

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
25 October 2001 (25.10.2001)

PCT

(10) International Publication Number
WO 01/79517 A2

(51) International Patent Classification⁷: C12N 15/82

(21) International Application Number: PCT/US01/11967

(22) International Filing Date: 13 April 2001 (13.04.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
09/548,971 13 April 2000 (13.04.2000) US

(71) Applicant: **THE REGENTS OF THE UNIVERSITY OF CALIFORNIA** [US/US]; University of California, San Diego Technology Transfer and Intellectual Property Services, 9500 Gilman Drive, MC 0910, La Jolla, CA 92093-0910 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:


— without international search report and to be republished upon receipt of that report

(72) Inventors: **LILJEGREN, Sarah**; 3268 Caminito Ameca, La Jolla, CA 92037 (US). **YANOFSKY, Martin, F.**; 5039 Manor Ridge Lane, San Diego, CA 92130 (US).

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(74) Agents: **HINSCH, Matthew, E.** et al.; Townsend and Townsend and Crew LLP, Two Embarcadero Center, Eighth Floor, San Francisco, CA 94111-3834 (US).

(54) Title: CONTROL OF FRUIT DEHISCENCE IN ARABIDOPSIS BY INDEHISCENT1 GENES



(57) Abstract: The present invention provides methods and compositions that modulate fruit dehiscence in plants.

WO 01/79517 A2

BEST AVAILABLE COPY

CONTROL OF FRUIT DEHISCENCE IN *ARABIDOPSIS* BY *INDEHISCENT1* GENES

5 CROSS-REFERENCES TO RELATED APPLICATIONS

This application is a continuation-in-part of U.S. Application Serial No. 09/339,998, filed on June 25, 1999, which claims benefit of priority to U.S. Provisional Application No. 60/090,649, filed June 25, 1998, each of which is incorporated by reference in its entirety. This application is also related to U.S. Application Serial No. 10 09/349,677, filed July 8, 1999, which is a divisional application of U.S. Application Serial No. 09/067,800, filed April 28, 1998, which claims the benefit of priority of U.S. Provisional Application No. 60/051,030, filed June 27, 1997, each of which is incorporated by reference in its entirety.

15 STATEMENT AS TO RIGHTS TO INVENTIONS MADE UNDER FEDERALLY SPONSORED RESEARCH AND DEVELOPMENT

This invention was made with Government support under National Science Foundation Grant number IBN-9985530. The Government has certain rights in the invention.

20 FIELD OF THE INVENTION

This invention relates plant genetic engineering. In particular, the invention relates to methods and compositions that modulate fruit dehiscence in plants.

BACKGROUND OF THE INVENTION

25 Rapeseed is one of the most important oilseed crops after soybeans and cottonseed, representing 10% of the world oilseed production in 1990. Rapeseed contains 40% oil, which is pressed from the seed, leaving a high-protein seed meal of value for animal feed and nitrogen fertilizer. Rapeseed oil, also known as canola oil, is a valuable product, representing the fourth most commonly traded vegetable oil in the world.

30 Unfortunately, the yield of seed from rapeseed and related plants is limited by pod dehiscence, which is a process that occurs late in fruit development whereby the pod is opened and the enclosed seeds released. Degradation and separation of cell walls along a discrete layer of cells dividing the two halves of the pod, termed the "dehiscence

zone," result in separation of the two halves of the pod and release of the contained seeds. The dehiscence zone is a region of only one to three cells in width that extends along the entire length of the valve/replum boundary (Meakin and Roberts, *J. Exp. Botany* 41:995-1002 (1990)). As the cells in the dehiscence zone separate from one another, the valves
5 detach from the replum, allowing seeds to be dispersed. Seed "shattering," whereby seeds are prematurely shed through dehiscence before the crop can be harvested, is a significant problem faced by commercial seed producers and represents a loss of income to the industry. Adverse weather conditions can exacerbate the process of dehiscence, resulting in greater than 50% loss of seed yield.

10 The fruit, a complex structure unique to flowering plants, mediates the maturation and dispersal of seeds. In most flowering plants, the fruit consists of the pericarp, which is derived from the ovary wall, and the seeds, which develop from fertilized ovules. *Arabidopsis*, which is typical of the more than 3000 species of the *Brassicaceae*, produces fruit in which the two carpel valves (ovary walls) are joined to
15 the replum, a visible suture that divides the two carpels.

 The plant hormone ethylene is produced by developing seeds and appears to be an important regulator of the dehiscence process. One line of evidence supporting a
role for ethylene in regulation of dehiscence comes from studies of fruit ripening, which, like fruit dehiscence, is a process involving the breakdown of cell wall material. In fruit
20 ripening, ethylene acts in part by activating cell wall degrading enzymes such as polygalacturonase (Theologis *et al.*, *Develop. Genetics* 14:282-295 (1993)). Moreover, in genetically modified tomato plants in which the ethylene response is blocked, such as transgenic tomato plants expressing antisense polygalacturonase, there is a significant delay in fruit ripening (Lanahan *et al.*, *The Plant Cell* 6:521-530 (1994); Smith *et al.*,
25 *Nature* 334:724-726 (1988)).

 In dehiscence, ultrastructural changes that culminate in degradation of the middle lamella of dehiscence zone cell walls weaken rapeseed pods and eventually lead to pod shatter. As in fruit ripening, hydrolytic enzymes including polygalacturonases play a role in this programmed breakdown. For example, in oilseed rape, a specific endo-
30 polygalacturonase, RDPG1, is upregulated and expressed exclusively in the dehiscence zone late in pod development (Petersen *et al.*, *Plant Mol. Biol.* 31:517-527 (1996), which is incorporated herein by reference). Ethylene may regulate the activity of hydrolytic enzymes involved in the process of dehiscence as it does in fruit ripening (Meakin and Roberts, *J. Exp. Botany* 41:1003-1011 (1990), which is incorporated herein by reference).

Yet, until now, the proteins that control the process of dehiscence, such as those regulating the relevant hydrolytic enzymes, have eluded identification.

Attempts to solve the problem of pod shatter and early fruit dehiscence over the past 20 years have focused on the breeding of shatter-resistant varieties.

5 However, these plant hybrids are frequently sterile and lose favorable characteristics that must be regained by backcrossing, which is both time-consuming and laborious. Other strategies to alleviate pod shattering include the use of chemicals such as pod sealants or mechanical techniques such as swathing to reduce wind-stimulated shattering. To date, however, a simple method for producing genetically modified plants that do not open and
10 release their seeds prematurely has not been described.

Thus, a need exists for identifying genes that regulate the dehiscence process and for developing genetically modified plant varieties in which the natural seed dispersal process is delayed. The present invention satisfies this need and provides related advantages as well.

15

SUMMARY OF THE INVENTION

This invention provides an isolated nucleic acid comprising an IND1 polynucleotide sequence encoding an IND1 polypeptide at least 70% identical to SEQ ID
20 NO:2. The isolated nucleic acid, for instance, can comprise a polynucleotide that encodes SEQ ID NO:2. In another embodiment, the isolated nucleic acid, for instance, can comprise positions from about 2765 to about 3361 of SEQ ID NO:1. For example, the nucleic acid can comprise SEQ ID NO:1.

The present invention also provides an expression cassette comprising a
25 promoter operably linked to an IND1 polynucleotide sequence, or complement thereof, encoding an IND1 polypeptide at least about 70% identical to SEQ ID NO:1. The expression cassette, for instance, can comprise a polynucleotide that encodes SEQ ID NO:2. In another embodiment, the expression cassette, for instance, can comprise positions from about 2765 to about 3361 of SEQ ID NO:1. For example, the expression
30 cassette can comprise SEQ ID NO:1. In some embodiments, the expression cassette comprises a promoter. The promoter, for instance, can be constitutive or tissue specific. In one aspect of the invention, the promoter is a dehiscence zone specific promoter. In another aspect, the promoter can comprise positions from about 1 to about 2764 or positions from about 3362 to about 3856 of SEQ ID NO:1.

The present invention also provides a plant comprising a recombinant expression cassette comprising a promoter operably linked to a polynucleotide sequence encoding an IND1 polypeptide at least about 70% identical to SEQ ID NO:1. In one aspect, the polynucleotide sequence encoding the IND1 polypeptide is operably linked to the promoter in the antisense orientation. In another aspect, the polynucleotide sequence encoding the IND1 polypeptide is operably linked to the promoter in the sense orientation. The polynucleotide sequence can further comprise a second polynucleotide sequence encoding the IND1 polypeptide wherein the second polynucleotide sequence is operably linked to a second promoter in the antisense orientation. In some embodiments, the plant of the invention has reduced lignification in valve margin cells. In other embodiments, the promoter is a dehiscence zone-selective regulatory element. In some of these embodiments, the regulatory element comprises positions from about 1 to about 2764 or from about 3362 to about 3856 of SEQ ID NO:1.

The present invention also provides a method of delaying fruit dehiscence in a plant comprising suppressing expression of an IND1 nucleic acid in the plant by introducing into the plant a recombinant expression cassette comprising a promoter operably linked to a polynucleotide sequence encoding an IND1 polypeptide at least 70% identical to SEQ ID NO:2. In some embodiments, the IND1 polypeptide is SEQ ID NO:2. In other embodiments, the IND1 polynucleotide comprises positions from about 2765 to about 3361 of SEQ ID NO:1. In one aspect of the invention, the IND1 polynucleotide comprises SEQ ID NO:1. The method can include a polynucleotide sequence encoding the IND1 polypeptide operably linked to the promoter in the antisense orientation. In another embodiment, the promoter is linked to the promoter in the sense orientation. The polynucleotide sequence can further comprise a second polynucleotide sequence encoding the IND1 polypeptide wherein the second polynucleotide sequence is operably linked to a second promoter in the antisense orientation. In some embodiments, the method results in a plant with reduced lignification in valve margin cells. In other embodiments, the promoter is a dehiscence zone-selective regulatory element. In some of these embodiments, the regulatory element comprises positions from about 1 to about 2764 or from about 3362 to about 3856 of SEQ ID NO:1. In one aspect, the recombinant expression cassette is introduced into the plant with *Agrobacterium*.

The present invention also provides a method of delaying fruit dehiscence in a plant comprising suppressing expression of an IND1 gene in the plant by introducing into the plant a recombinant expression cassette comprising a polynucleotide sequence at

least about 70% identical to positions from about 1 to about 2764 or from about 3362 to about 3856 of SEQ ID NO:1. In one aspect of the invention, the polynucleotide sequence comprises positions from about 1 to about 2764 or from about 3362 to about 3856 of SEQ ID NO:1. In some aspects, lignification is reduced in valve margin cells.

5

DEFINITIONS

The phrase "nucleic acid" refers to a single or double-stranded polymer of deoxyribonucleotide or ribonucleotide bases read from the 5' to the 3' end. Nucleic acids may also include modified nucleotides that permit correct read through by a polymerase and do not alter expression of a polypeptide encoded by that nucleic acid.

10

The phrase "polynucleotide sequence" or "nucleic acid sequence" includes both the sense and antisense strands of a nucleic acid as either individual single strands or in the duplex. It includes, but is not limited to, self-replicating plasmids, chromosomal sequences, and infectious polymers of DNA or RNA.

15

The phrase "nucleic acid sequence encoding" refers to a nucleic acid which directs the expression of a specific protein or peptide. The nucleic acid sequences include both the DNA strand sequence that is transcribed into RNA and the RNA sequence that is translated into protein. The nucleic acid sequences include both the full length nucleic acid sequences as well as non-full length sequences derived from the full length sequences. It should be further understood that the sequence includes the degenerate codons of the native sequence or sequences which may be introduced to provide codon preference in a specific host cell.

20

The term "promoter" or "regulatory element" refers to a region or sequence determinants located upstream or downstream from the start of transcription and which are involved in recognition and binding of RNA polymerase and other proteins to initiate transcription. A "plant promoter" is a promoter capable of initiating transcription in plant cells. Such promoters need not be of plant origin, for example, promoters derived from plant viruses, such as the CaMV35S promoter, can be used in the present invention.

25

As used herein, the term "dehiscence zone-selective regulatory element" refers to a nucleotide sequence that, when operatively linked to a nucleic acid molecule, confers selective expression upon the operatively linked nucleic acid molecule in a limited number of plant tissues, including the valve margin or dehiscence zone. The valve margin is the future site of the dehiscence zone and encompasses the margins of the outer replum as well as valve cells adjacent to the outer replum. The dehiscence zone,

30

which develops in the region of the valve margin, refers to the group of cells that separate during the process of dehiscence, allowing valves to come apart from the replum and the enclosed seeds to be released. Thus, a dehiscence zone-selective regulatory element, as defined herein, confers selective expression in the mature dehiscence zone, or confers
5 selective expression in the valve margin, which marks the future site of the dehiscence zone.-

A dehiscence zone-selective regulatory element can confer specific expression exclusively in cells of the valve margin or dehiscence zone or can confer selective expression in a limited number of plant cell types including cells of the valve
10 margin or dehiscence zone. A *SHATTERPROOF1* or *SHATTERPROOF2* (*SHP1* and *SHP2*, also designated as *AGL1* and *AGL5*, respectively) regulatory element, for example, which confers selective expression in ovules and placenta as well as in the dehiscence zone, is a dehiscence zone-selective regulatory element as defined herein. Similarly, an
15 *IND1* regulatory element also confers selective expression in the dehiscence zone. A dehiscence zone-selective regulatory element generally is distinguished from other regulatory elements by conferring selective expression in the valve margin or dehiscence zone without conferring expression throughout the adjacent carpel valves.

It is understood that limited modifications can be made without destroying the biological function of a regulatory element and that such limited modifications can
20 result in dehiscence zone-selective regulatory elements that have substantially equivalent or enhanced function as compared to a wild type *IND1* regulatory element. These modifications can be deliberate, as through site-directed mutagenesis, or can be accidental such as through mutation in hosts harboring the regulatory element. All such modified nucleotide sequences are included in the definition of a dehiscence zone-selective
25 regulatory element as long as the ability to confer selective expression in the valve margin or dehiscence zone is substantially retained.

The term "plant" includes whole plants, shoot vegetative organs/structures (e.g. leaves, stems and tubers), roots, flowers and floral organs/structures (e.g. bracts, sepals, petals, stamens, carpels, anthers and ovules), seed (including embryo, endosperm,
30 and seed coat) and fruit (the mature ovary), plant tissue (e.g. vascular tissue, ground tissue, and the like) and cells (e.g. guard cells, egg cells, trichomes and the like), and progeny of same. The class of plants that can be used in the method of the invention is generally as broad as the class of higher and lower plants amenable to transformation techniques, including angiosperms (monocotyledonous and dicotyledonous plants),

gymnosperms, ferns, and multicellular algae. It includes plants of a variety of ploidy levels, including aneuploid, polyploid, diploid, haploid and hemizygous.

The term "seed plant" means an angiosperm or gymnosperm. An angiosperm is a seed-bearing plant whose seeds are borne in a mature ovary (fruit). An angiosperm commonly is recognized as a flowering plant. Angiosperms are divided into two broad classes based on the number of cotyledons, which are seed leaves that generally store or absorb food. Thus, a monocotyledonous angiosperm is an angiosperm having a single cotyledon, whereas a dicotyledonous angiosperm is an angiosperm having two cotyledons. A variety of angiosperms are known including, for example, oilseed plants, leguminous plants, fruit-bearing plants, ornamental flowers, cereal plants and hardwood trees, which general classes are not necessarily exclusive. The skilled artisan will recognize that the methods of the invention can be practiced using these or other angiosperms, as desired. A gymnosperm is a seed-bearing plant with seeds not enclosed in an ovary.

The phrase "host cell" refers to a cell from any organism. Preferred host cells are derived from plants, bacteria, yeast, fungi, insects or other animals. Methods for introducing polynucleotide sequences into various types of host cells are well known in the art.

The term "delayed," as used herein in reference to the timing of seed dispersal in a fruit produced by a non-naturally occurring plant of the invention, means a significantly later time of seed dispersal as compared to the time seeds normally are dispersed from a corresponding plant at the same developmental stage expressing naturally-occurring levels of IND1. Thus, the term "delayed" is used broadly to encompass both seed dispersal that is significantly postponed as compared to the seed dispersal in a corresponding plant, and to seed dispersal that is completely precluded, such that fruits never release their seeds unless there is human or other intervention.

It is recognized that there can be natural variation of the time of seed dispersal within a plant species or variety. However, a "delay" in the time of seed dispersal in a non-naturally occurring plant of the invention readily can be identified by sampling a population of the non-naturally occurring plants and determining that the normal distribution of seed dispersal times is significantly later, on average, than the normal distribution of seed dispersal times in a population of the corresponding plant species or variety that does not contain an exogenous IND1 polynucleotide. Thus, production of non-naturally occurring plants of the invention provides a means to skew

the normal distribution of the time of seed dispersal from pollination, such that seeds are dispersed, on average, at least about 1%, 2%, 5%, 10%, 30%, 50%, 100%, 200% or 500% later than in the corresponding plant species that does not contain an exogenous nucleic acid molecule encoding an *IND1* gene product.

5 The term "suppressed" or "decreased" encompasses the absence of *IND1* protein in a plant, as well as protein expression that is present but reduced as compared to the level of *IND1* protein expression in a wild type plant. Furthermore, the term suppressed refers to *IND1* protein expression that is reduced throughout the entire domain of *IND1* expression, or to expression that is reduced in some part of the *IND1* expression
10 domain, provided that the resulting plant is characterized by delayed seed dispersal. The term "suppressed" also encompasses an amount of *IND1* protein that is equivalent to wild type *IND1* expression, but where the *IND1* protein has a reduced level of activity. As discussed above, *IND1* each contain a conserved an basic HLH domain; point mutations or gross deletions within the HLH domain that reduce the DNA-binding activity of *IND1*
15 can reduce or destroy the activity of *IND1* and, therefore, "suppress" *IND1* expression as defined herein. One skilled in the art will recognize that, preferably, *IND1* expression is essentially absent in the valve margin of a plant or the *IND1* protein is essentially non-functional.

 "Increased" or "enhanced" *IND1* activity or expression of a *IND1* gene
20 refers to an augmented change in *IND1* activity. Examples of such increased activity or expression include the following. *IND1* activity or expression of the *IND1* gene is increased above the level of that in wild-type, non-transgenic control plants (i.e. the quantity of *IND1* activity or expression of the *IND1* gene is increased). *IND1* activity or expression of the *IND1* gene is in an organ, tissue or cell where it is not normally detected
25 in wild-type, non-transgenic control plants (i.e. spatial distribution of *IND1* activity or expression of the *IND1* gene is increased). *IND1* activity or expression is increased when *IND1* activity or expression of the *IND1* gene is present in an organ, tissue or cell for a longer period than in a wild-type, non-transgenic controls (i.e. duration of *IND1* activity or expression of the *IND1* gene is increased).

30 A polynucleotide sequence is "heterologous to" a second polynucleotide sequence if it originates from a foreign species, or, if from the same species, is modified by human action from its original form. For example, a promoter operably linked to a heterologous coding sequence refers to a coding sequence from a species different from

that from which the promoter was derived, or, if from the same species, a coding sequence which is different from any naturally occurring allelic variants.

A polynucleotide "exogenous to" an individual plant is a polynucleotide which is introduced into the plant, or a predecessor generation of the plant, by any means other than by a sexual cross. An exogenous nucleic acid molecule can have a naturally occurring or non-naturally occurring nucleotide sequence and can be a heterologous nucleic acid molecule derived from a different plant species than the plant into which the nucleic acid molecule is introduced or can be a nucleic acid molecule derived from the same plant species as the plant into which it is introduced. Examples of means by which this can be accomplished are described below, and include *Agrobacterium*-mediated transformation, biolistic methods, electroporation, in planta techniques, and the like.

An "*IND1* polynucleotide" is a nucleic acid sequence comprising (or consisting of) a coding region of about 50 to about 4000 nucleotides, sometimes from about 100 to about 3000 nucleotides and sometimes from about 200 to about 600 nucleotides, which hybridizes to SEQ ID NO:1 under stringent conditions (as defined below), or which encodes an *IND1* polypeptide or fragment of at least 15 amino acids thereof. *IND1* polynucleotides can also be identified by their ability to hybridize under low stringency conditions (e.g., $T_m \sim 40^\circ\text{C}$) to nucleic acid probes having the sequence of SEQ ID NO:1. SEQ ID NO:1 is an example of a *IND1* polynucleotide.

A "promoter from a *IND1* gene" or "*IND1* promoter" will typically be about 500 to about 3000 nucleotides in length, usually from about 750 to 2750. Exemplary promoter sequences are shown as SEQ ID NO:3 and SEQ ID NO:4. SEQ ID NO:3 represents the 5' untranslated region of the *IND1* and SEQ ID NO:4 represents the 3' untranslated region of *IND1*. A *IND1* promoter can also be identified by its ability to direct expression in the valve margin of fruit. In particular, the *Ind1* promoter directs expression at the valve margin of developing gynoecium just prior to fertilization (stage 13) through the maturation of the fruit (stage 17). The promoter does not provide significant expression in leaf tissue.

An "*IND1* polypeptide" is a sequence of about 50 to about 200, sometimes 100 to 190, and preferably 198 amino acid residues encoded by a *IND1* polynucleotide. *IND1* polypeptides are characterized by the presence of an basic helix-loop-helix (HLH) domain which bind specific polynucleotide sequences. For instance amino acid residues ISDDPQTVVARRRRERISEKIRILKRIVPGGAKMDTASMLDEAIRYTKFLK represent the HLH domain of the polypeptide shown in SEQ ID NO:2. The HLH domain

is known in the art and is shared by other transcription factors including uncharacterized sequences represented by GenBank accession number E1283552 and 2262147 and the gene product, PIF3 (Ni *et al. Cell* 95:657 (1998)). The HLH domain of IND1 is therefore a DNA binding domain.

5 As used herein, a homolog of a particular IND1 gene (e.g., SEQ ID NO:1) is a second gene in the same plant type or in a different plant type, which has a polynucleotide sequence of at least 50 contiguous nucleotides which are substantially identical (determined as described below) to a sequence in the first gene. It is believed that, in general, homologs share a common evolutionary past.

10 A "polynucleotide sequence from" a particular gene is a subsequence or full length polynucleotide sequence of an IND1 gene which, when present in a transgenic plant, has the desired effect. For example, one effect is inhibition of expression of the endogenous gene driving expression of an heterologous polynucleotide. A full length sequence of a particular gene disclosed here may contain about 95%, usually at least
15 about 98% of an entire sequence shown in the Sequence Listing, below.

 The term "reproductive tissues" as used herein includes fruit, ovules, seeds, pollen, pistils, flowers, or any embryonic tissue.

 An "expression cassette" refers to a nucleic acid construct, which when introduced into a host cell, results in transcription and/or translation of a RNA or
20 polypeptide, respectively. Antisense or sense constructs that are not or cannot be translated are expressly included by this definition.

 In the case of both expression of transgenes and inhibition of endogenous genes (e.g., by antisense, or sense suppression) one of skill will recognize that the inserted polynucleotide sequence need not be identical and may be "substantially identical" to a
25 sequence of the gene from which it was derived. As explained below, these variants are specifically covered by this term.

 In the case where the inserted polynucleotide sequence is transcribed and translated to produce a functional polypeptide, one of skill will recognize that because of codon degeneracy a number of polynucleotide sequences will encode the same
30 polypeptide. These variants are specifically covered by the term "polynucleotide sequence from" a particular valve-margin gene, such as *IND1*. In addition, the term specifically includes sequences (e.g., full length sequences) substantially identical (determined as described below) with a IND1 gene sequence and that encode proteins that retain the function of a IND1 polypeptide.

In the case of polynucleotides used to inhibit expression of an endogenous gene, the introduced sequence need not be perfectly identical to a sequence of the target endogenous gene. The introduced polynucleotide sequence will typically be at least substantially identical (as determined below) to the target endogenous sequence.

5 Two nucleic acid sequences or polypeptides are said to be "identical" if the sequence of nucleotides or amino acid residues, respectively, in the two sequences is the same when aligned for maximum correspondence as described below. The term "complementary to" is used herein to mean that the sequence is complementary to all or a portion of a reference polynucleotide sequence.

10 Optimal alignment of sequences for comparison may be conducted by the local homology algorithm of Smith and Waterman *Add. APL. Math.* 2:482 (1981), by the homology alignment algorithm of Needleman and Wunsch *J. Mol. Biol.* 48:443 (1970), by the search for similarity method of Pearson and Lipman *Proc. Natl. Acad. Sci. (U.S.A.)* 85: 2444 (1988), by computerized implementations of these algorithms (GAP, BESTFIT, 15 BLAST, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group (GCG), 575 Science Dr., Madison, WI), or by inspection.

"Percentage of sequence identity" is determined by comparing two optimally aligned sequences over a comparison window, wherein the portion of the polynucleotide sequence in the comparison window may comprise additions or deletions 20 (i.e., gaps) as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. The percentage is calculated by determining the number of positions at which the identical nucleic acid base or amino acid residue occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of 25 comparison and multiplying the result by 100 to yield the percentage of sequence identity.

The term "substantial identity" of polynucleotide sequences means that a polynucleotide comprises a sequence that has at least 25% sequence identity. Alternatively, percent identity can be any integer from 25% to 100%. More preferred 30 embodiments include at least: 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, or 99%. compared to a reference sequence using the programs described herein; preferably BLAST using standard parameters, as described below. Accordingly, IND1 sequences of the invention include nucleic acid sequences that have substantial identity to SEQ ID NO:1. IND1 sequences of the invention also include polypeptide sequences having substantial identity to SEQ ID NO:2. One of skill

will recognize that these values can be appropriately adjusted to determine corresponding identity of proteins encoded by two nucleotide sequences by taking into account codon degeneracy, amino acid similarity, reading frame positioning and the like. Substantial identity of amino acid sequences for these purposes normally means sequence identity of

5 at least 40%. Preferred percent identity of polypeptides can be any integer from 40% to 100%. More preferred embodiments include at least 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, or 99%. Most preferred embodiments include 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74% and 75%.

Polypeptides which are "substantially similar" share sequences as noted above except that

10 residue positions which are not identical may differ by conservative amino acid changes. Conservative amino acid substitutions refer to the interchangeability of residues having similar side chains. For example, a group of amino acids having aliphatic side chains is glycine, alanine, valine, leucine, and isoleucine; a group of amino acids having aliphatic-hydroxyl side chains is serine and threonine; a group of amino acids having amide-

15 containing side chains is asparagine and glutamine; a group of amino acids having aromatic side chains is phenylalanine, tyrosine, and tryptophan; a group of amino acids having basic side chains is lysine, arginine, and histidine; and a group of amino acids having sulfur-containing side chains is cysteine and methionine. Preferred conservative amino acids substitution groups are: valine-leucine-isoleucine, phenylalanine-tyrosine,

20 lysine-arginine, alanine-valine, aspartic acid-glutamic acid, and asparagine-glutamine.

Another indication that nucleotide sequences are substantially identical is if two molecules hybridize to each other, or a third nucleic acid, under stringent conditions. Stringent conditions are sequence dependent and will be different in different circumstances. Generally, stringent conditions are selected to be about 5°C lower than

25 the thermal melting point (T_m) for the specific sequence at a defined ionic strength and pH. The T_m is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. Typically, stringent conditions will be those in which the salt concentration is about 0.02 molar at pH 7 and the temperature is at least about 60°C.

30 In the present invention, mRNA encoded by IND1 genes of the invention can be identified in Northern blots under stringent conditions using cDNAs of the invention or fragments of at least about 100 nucleotides. For the purposes of this disclosure, stringent conditions for such RNA-DNA hybridizations are those which include at least one wash in 0.2X SSC at 63°C for 20 minutes, or equivalent conditions.

Genomic DNA or cDNA comprising genes of the invention can be identified using the same cDNAs (or fragments of at least about 100 nucleotides) under stringent conditions, which for purposes of this disclosure, include at least one wash (usually 2) in 0.2X SSC at a temperature of at least about 50°C, usually about 55°C, for 20 minutes, or equivalent
5 conditions.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 illustrates the cell types of the Arabidopsis fruit at maturity.

10

DETAILED DESCRIPTION

I. Introduction

The present invention provides methods of modulating fruit development in plants. In particular, the invention provides methods of delaying or preventing fruit
15 dehiscence by suppressing expression of an bHLH gene such as IND1 in a plant. The invention also provides transgenic plants comprising various polynucleotides encoding an bHLH polypeptide such as IND1.

The present invention relates to the previous discovery that an *agl1 agl2* double mutant plant has a delayed seed dispersal phenotype (Liljegren *et al.*, *Nature*
20 404:766-770 (2000)). Loss-of-function mutations in the *SHP1* and *SHP2* genes were produced by disruptive T-DNA insertion and homologous recombination. In the resulting *shp1 shp2* double mutant plants, the dehiscence zone failed to develop normally, and the mature fruits did not undergo dehiscence. Thus, *SHP1* or *SHP2* gene expression is required for development of the dehiscence zone. These results indicate that *SHP1* and
25 *SHP2* regulate pod dehiscence and that manipulation of *SHP1* and *SHP2* expression can allow the process of pod shatter to be controlled.

The present invention provides evidence that *IND1* is regulated by *SHP1* and *SHP2* and that expression of *IND1* modulates fruit dehiscence. The present invention also provides for methods of delaying fruit dehiscence by suppressing expression of
30 *IND1*.

The Arabidopsis *SHP1* and *SHP2* genes encode MADS box proteins with 85% identity at the amino acid level. The *SHP1* and *SHP2* RNA expression patterns are also strikingly similar. In particular, both RNAs are specifically expressed in flowers, where they accumulate in developing carpels. In particular, strong expression of these

genes is observed in the outer replum along the valve/replum boundary (Ma *et al.*, *supra*, 1991; Savidge *et al.*, *The Plant Cell* 7:721-723 (1995); Flanagan *et al.*, *The Plant Journal* 10:343-353 (1996), each of which is incorporated herein by reference). Thus, *SHP1* and *SHP2* are expressed in the valve margin, at least within the cells of the outer replum.

5 Generally, the nomenclature and the laboratory procedures in recombinant DNA technology described below are those well known and commonly employed in the art. Standard techniques are used for cloning, DNA and RNA isolation, amplification and purification. Generally enzymatic reactions involving DNA ligase, DNA polymerase, restriction endonucleases and the like are performed according to the manufacturer's
10 specifications. These techniques and various other techniques are generally performed according to Sambrook *et al.*, *Molecular Cloning - A Laboratory Manual*, 2nd. ed., Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, (1989).

II. Isolation of nucleic acids of the invention

15 The isolation of sequences from the genes of the invention may be accomplished by a number of techniques. For instance, oligonucleotide probes based on the sequences disclosed here can be used to identify the desired gene in a cDNA or genomic DNA library from a desired plant species. To construct genomic libraries, large segments of genomic DNA are generated by random fragmentation, e.g. using restriction
20 endonucleases, and are ligated with vector DNA to form concatemers that can be packaged into the appropriate vector. To prepare a library of embryo-specific cDNAs such as *IND1*, mRNA is isolated from embryos and a cDNA library that contains the gene transcripts is prepared from the mRNA.

 The cDNA or genomic library can then be screened using a probe based
25 upon the sequence of a cloned *IND1* gene such as the polynucleotides disclosed here. Probes may be used to hybridize with genomic DNA or cDNA sequences to isolate homologous genes in the same or different plant species.

 Alternatively, the nucleic acids of interest can be amplified from nucleic acid samples using amplification techniques. For instance, polymerase chain reaction
30 (PCR) technology to amplify the sequences of the genes directly from mRNA, from cDNA, from genomic libraries or cDNA libraries. PCR and other *in vitro* amplification methods may also be useful, for example, to clone nucleic acid sequences that code for proteins to be expressed, to make nucleic acids to use as probes for detecting the presence of the desired mRNA in samples, for nucleic acid sequencing, or for other purposes.

Appropriate primers and probes for identifying genes such as *IND1* from plant tissues are generated from comparisons of the sequences provided herein. For a general overview of PCR see PCR Protocols: A Guide to Methods and Applications. (Innis, M, Gelfand, D., Sninsky, J. and White, T., eds.), Academic Press, San Diego

5 (1990). Appropriate primers for amplification of the genomic region of *IND1* or the *IND1* cDNA include the following primer pairs: 5'-gatgaaatggaaatggtatgtata-3' and 5'-gttcacagggttgggagttgtg-3'. The amplification conditions are typically as follows. Reaction components: 10 mM Tris-HCl, pH 8.3, 50 mM potassium chloride, 1.5 mM magnesium chloride, 0.001% gelatin, 200 μ M dATP, 200 μ M dCTP, 200 μ M dGTP, 200

10 μ M dTTP, 0.4 μ M primers, and 100 units per ml Taq polymerase. Program: 96 C for 3 min., 30 cycles of 96 C for 45 sec., 50 C for 60 sec., 72 for 60 sec, followed by 72 C for 5 min.

Polynucleotides may also be synthesized by well-known techniques as described in the technical literature. See, e.g., Carruthers *et al.*, Cold Spring Harbor

15 Symp. *Quant. Biol.* 47:411-418 (1982), and Adams *et al.*, *J. Am. Chem. Soc.* 105:661 (1983). Double stranded DNA fragments may then be obtained either by synthesizing the complementary strand and annealing the strands together under appropriate conditions, or by adding the complementary strand using DNA polymerase with an appropriate primer sequence.

20 The genus of *IND1* nucleic acid sequences of the invention includes genes and gene products identified and characterized by analysis using nucleic acid sequences of the invention, including SEQ ID NO:1 and protein sequences of the invention, including SEQ ID NO:2. *IND1* sequences of the invention include nucleic acid sequences having substantial identity to SEQ ID NO:1. *IND1* sequences of the invention

25 also include polypeptide sequences having substantial identity to SEQ ID NO:2.

III. Use of nucleic acids of the invention

30 A. Use of nucleic acids of the invention to inhibit or suppress gene expression

The invention provides methods of modulating fruit dehiscence in a plant by introducing into a plant a recombinant expression cassette comprising a regulatory element operably linked to a HLH polynucleotide such as *IND1*. The invention also provides methods for delaying seed dispersal in a plant by suppressing expression of a

nucleic acid molecule encoding an IND1 gene product. In a transgenic plant of the invention, a nucleic acid molecule, or antisense constructs thereof, encoding an IND1 gene product can be operatively linked to an exogenous regulatory element. The invention provides, for example, a transgenic plant characterized by delayed seed dispersal having an expressed nucleic acid molecule encoding an IND1 gene product, or antisense construct thereof, that is operatively linked to an exogenous constitutive regulatory element. In one embodiment, the invention provides a transgenic plant that is characterized by delayed seed dispersal due to suppression of a nucleic acid molecule encoding an IND1 ortholog. In some preferred embodiments, suppression of IND1 expression results in reduced lignification in valve margin cells. *See also*, U.S. Application Serial No. 09/339,998, filed on June 25, 1999.

The *IND1* sequences prepared as described herein, can be used to prepare expression cassettes useful in a number of techniques, including inhibiting or suppressing expression. A number of methods can be used to inhibit gene expression in plants. For instance, antisense technology can be conveniently used. To accomplish this, a nucleic acid segment from the desired gene is cloned and operably linked to a promoter such that the antisense strand of RNA will be transcribed. The expression cassette is then transformed into plants and the antisense strand of RNA is produced. In plant cells, it has been suggested that antisense RNA inhibits gene expression by preventing the accumulation of mRNA which encodes the enzyme of interest, see, e.g., Sheehy *et al.*, *Proc. Nat. Acad. Sci. USA*, 85:8805-8809 (1988); Pnueli *et al.*, *The Plant Cell* 6:175-186 (1994); and Hiatt *et al.*, U.S. Patent No. 4,801,340.

The antisense nucleic acid sequence transformed into plants will be substantially identical to at least a portion of the endogenous gene or genes to be repressed. The sequence, however, does not have to be perfectly identical to inhibit expression. Thus, an antisense or sense nucleic acid molecule encoding only a portion of IND1 can be useful for producing a plant in which IND1 expression is suppressed. The vectors of the present invention can be designed such that the inhibitory effect applies to other proteins within a family of genes exhibiting homology or substantial homology to the target gene.

For antisense suppression, the introduced sequence also need not be full length relative to either the primary transcription product or fully processed mRNA. Generally, higher homology can be used to compensate for the use of a shorter sequence. Furthermore, the introduced sequence need not have the same intron or exon pattern, and

homology of non-coding segments may be equally effective. Normally, a sequence of between about 30 or 40 nucleotides and about full length nucleotides should be used, though a sequence of at least about 100 nucleotides is preferred, a sequence of at least about 200 nucleotides is more preferred, and a sequence of at least about 500 nucleotides is especially preferred.

Catalytic RNA molecules or ribozymes can also be used to inhibit expression of *IND1* genes. It is possible to design ribozymes that specifically pair with virtually any target RNA and cleave the phosphodiester backbone at a specific location, thereby functionally inactivating the target RNA. In carrying out this cleavage, the ribozyme is not itself altered, and is thus capable of recycling and cleaving other molecules, making it a true enzyme. The inclusion of ribozyme sequences within antisense RNAs confers RNA-cleaving activity upon them, thereby increasing the activity of the constructs.

A number of classes of ribozymes have been identified. One class of ribozymes is derived from a number of small circular RNAs that are capable of self-cleavage and replication in plants. The RNAs replicate either alone (viroid RNAs) or with a helper virus (satellite RNAs). Examples include RNAs from avocado sunblotch viroid and the satellite RNAs from tobacco ringspot virus, lucerne transient streak virus, velvet tobacco mottle virus, solanum nodiflorum mottle virus and subterranean clover mottle virus. The design and use of target RNA-specific ribozymes is described in Haseloff *et al. Nature*, 334:585-591 (1988).

Another method of suppression is sense suppression (also known as co-suppression). Introduction of expression cassettes in which a nucleic acid is configured in the sense orientation with respect to the promoter has been shown to be an effective means by which to block the transcription of target genes. For an example of the use of this method to modulate expression of endogenous genes see, Napoli *et al.*, *The Plant Cell* 2:279-289 (1990); Flavell, *Proc. Natl. Acad. Sci., USA* 91:3490-3496 (1994); Kooter and Mol, *Current Opin. Biol.* 4:166-171 (1993); and U.S. Patents Nos. 5,034,323, 5,231,020, and 5,283,184.

Generally, where inhibition of expression is desired, some transcription of the introduced sequence occurs. The effect may occur where the introduced sequence contains no coding sequence per se, but only intron or untranslated sequences homologous to sequences present in the primary transcript of the endogenous sequence. The introduced sequence generally will be substantially identical to the endogenous

sequence intended to be repressed. This minimal identity will typically be greater than about 65%, but a higher identity might exert a more effective repression of expression of the endogenous sequences. Substantially greater identity of more than about 80% is preferred, though about 95% to absolute identity would be most preferred. As with
5 antisense regulation, the effect should apply to any other proteins within a similar family of genes exhibiting homology or substantial homology.

For sense suppression, the introduced sequence in the expression cassette, needing less than absolute identity, also need not be full length, relative to either the primary transcription product or fully processed mRNA. This may be preferred to avoid
10 concurrent production of some plants that are overexpressers. A higher identity in a shorter than full length sequence compensates for a longer, less identical sequence. Furthermore, the introduced sequence need not have the same intron or exon pattern, and identity of non-coding segments will be equally effective. Normally, a sequence of the size ranges noted above for antisense regulation is used.

15 In a preferred embodiment, expression of a nucleic acid of interest can be suppressed by the simultaneous expression of both sense and antisense constructs (Waterhouse *et al.*, *Proc. Natl. Acad. Sci. USA* 95:13959-13964 (1998). See also Tabara *et al. Science* 282:430-431 (1998).

One of skill in the art will recognize that using technology based on
20 specific nucleotide sequences (e.g., antisense or sense suppression technology), families of homologous genes can be suppressed with a single sense or antisense transcript. For instance, if a sense or antisense transcript is designed to have a sequence that is conserved among a family of genes, then multiple members of a gene family can be suppressed. Conversely, if the goal is to only suppress one member of a homologous gene family,
25 then the sense or antisense transcript should be targeted to sequences with the most variance between family members.

Another means of inhibiting *IND1* function in a plant is by creation of dominant negative mutations. In this approach, non-functional, mutant *IND1* polypeptides, which retain the ability to interact with wild-type subunits are introduced
30 into a plant. A dominant negative construct also can be used to suppress *IND1* expression in a plant. A dominant negative construct useful in the invention generally contains a portion of the complete *IND1* coding sequence sufficient, for example, for DNA-binding or for a protein-protein interaction such as a homodimeric or heterodimeric protein-protein interaction but lacking the transcriptional activity of the wild type protein. For

example, a carboxy-terminal deletion mutant of AGAMOUS was used as a dominant negative construct to suppress expression of the MADS box gene AGAMOUS (Mizukami et al., Plant Cell 8:831-844 (1996)). One skilled in the art understands that, similarly, a dominant negative IND1 construct can be used to suppress IND1 expression
5 in a plant.

B. Use of nucleic acids of the invention to enhance gene expression

Isolated sequences prepared as described herein can also be used to prepare expression cassettes that enhance or increase endogenous *IND1* gene expression.
10 Where overexpression of a gene is desired, the desired gene from a different species may be used to decrease potential sense suppression effects. Enhanced expression of *IND1* polynucleotides is useful, for example, to produce plants with small fruit.

Any of a number of means well known in the art can be used to increase *IND1* activity in plants. Any organ can be targeted, such as shoot vegetative
15 organs/structures (e.g. leaves, stems and tubers), roots, flowers and floral organs/structures (e.g. bracts, sepals, petals, stamens, carpels, anthers and ovules), seed (including embryo, endosperm, and seed coat) and fruit. Alternatively, one or several *IND1* genes can be expressed constitutively (e.g., using the CaMV 35S promoter).

One of skill will recognize that the polypeptides encoded by the genes of
20 the invention, like other proteins, have different domains which perform different functions. Thus, the gene sequences need not be full length, so long as the desired functional domain of the protein is expressed. As explained above, *IND1* polypeptides carry a bHLH domain, which is capable of binding DNA. Thus, without being bound to any particular theory or mechanism, *IND1* is likely to act as a transcriptional modulator.

25

C. Modification of endogenous IND1 genes

Methods for introducing genetic mutations into plant genes and selecting plants with desired traits are well known. For instance, seeds or other plant material can be treated with a mutagenic chemical substance, according to standard techniques. Such
30 chemical substances include, but are not limited to, the following: diethyl sulfate, ethylene imine, ethyl methanesulfonate and N-nitroso-N-ethylurea. Alternatively, ionizing radiation from sources such as, X-rays or gamma rays can be used.

Modified protein chains can also be readily designed utilizing various recombinant DNA techniques well known to those skilled in the art and described for

instance, in Sambrook et al., supra. Hydroxylamine can also be used to introduce single base mutations into the coding region of the gene (Sikorski, *et al.*, (1991). *Meth. Enzymol.* 194: 302-318). For example, the chains can vary from the naturally occurring sequence at the primary structure level by amino acid substitutions, additions, deletions, and the like.

- 5 These modifications can be used in a number of combinations to produce the final modified protein chain.

Alternatively, homologous recombination can be used to induce targeted gene modifications by specifically targeting the *IND1* gene *in vivo* (*see, generally*, Grewal and Klar, *Genetics* 146: 1221-1238 (1997) and Xu *et al.*, *Genes Dev.* 10: 2411-2422 (1996)). Homologous recombination has been demonstrated in plants (Puchta *et al.*, *Experientia* 50: 277-284 (1994), Swoboda *et al.*, *EMBO J.* 13: 484-489 (1994); Offringa *et al.*, *Proc. Natl. Acad. Sci. USA* 90: 7346-7350 (1993); and Kempin *et al.* *Nature* 389:802-803 (1997)).

In applying homologous recombination technology to the genes of the invention, mutations in selected portions of an *IND1* gene sequences (including 5' upstream, 3' downstream, and intragenic regions) such as those disclosed here are made *in vitro* and then introduced into the desired plant using standard techniques. Since the efficiency of homologous recombination is known to be dependent on the vectors used, use of dicistronic gene targeting vectors as described by Mountford *et al.*, *Proc. Natl. Acad. Sci. USA* 91: 4303-4307 (1994); and Vaultont *et al.*, *Transgenic Res.* 4: 247-255 (1995) are conveniently used to increase the efficiency of selecting for altered *IND1* gene expression in transgenic plants. The mutated gene will interact with the target wild-type gene in such a way that homologous recombination and targeted replacement of the wild-type gene will occur in transgenic plant cells, resulting in suppression of *IND1* activity.

Alternatively, oligonucleotides composed of a contiguous stretch of RNA and DNA residues in a duplex conformation with double hairpin caps on the ends can be used. The RNA/DNA sequence is designed to align with the sequence of the target *IND1* gene and to contain the desired nucleotide change. Introduction of the chimeric oligonucleotide on an extrachromosomal T-DNA plasmid results in efficient and specific *IND1* gene conversion directed by chimeric molecules in a small number of transformed plant cells. This method is described in Cole-Strauss *et al.*, *Science* 273:1386-1389 (1996) and Yoon *et al.*, *Proc. Natl. Acad. Sci. USA* 93: 2071-2076 (1996).

In other embodiments, the promoters derived from the *IND1* genes of the invention can be used to drive expression of heterologous genes in an valve margin-

specific manner. Suitable structural genes that could be used for this purpose include genes encoding cytotoxic proteins as discussed below.

Typically, desired promoters are identified by analyzing the 5' sequences of a genomic clone corresponding to the IND1 genes described here. Sequences characteristic of promoter sequences can be used to identify the promoter. Sequences controlling eukaryotic gene expression have been extensively studied. For instance, promoter sequence elements include the TATA box consensus sequence (TATAAT), which is usually 20 to 30 base pairs upstream of the transcription start site. In most instances the TATA box is required for accurate transcription initiation. In plants, further upstream from the TATA box, at positions -80 to -100, there is typically a promoter element with a series of adenines surrounding the trinucleotide G (or T) N G. J. Messing *et al.*, in GENETIC ENGINEERING IN PLANTS, pp.221-227 (Kosage, Meredith and Hollaender, eds. (1983)).

A number of methods are known to those of skill in the art for identifying and characterizing promoter regions in plant genomic DNA (see, e.g., Jordano, *et al.*, *Plant Cell*, 1: 855-866 (1989); Bustos, *et al.*, *Plant Cell*, 1:839-854 (1989); Green, *et al.*, *EMBO J.* 7, 4035-4044 (1988); Meier, *et al.*, *Plant Cell*, 3, 309-316 (1991); and Zhang, *et al.*, *Plant Physiology* 110: 1069-1079 (1996)).

IV. Preparation of recombinant vectors

To use isolated sequences in the above techniques, recombinant DNA vectors suitable for transformation of plant cells are prepared. Techniques for transforming a wide variety of higher plant species are well known and described in the technical and scientific literature. See, for example, Weising *et al. Ann. Rev. Genet.* 22:421-477 (1988). A DNA sequence coding for the desired polypeptide, for example a cDNA sequence encoding a full length protein, will preferably be combined with transcriptional and translational initiation regulatory sequences which will direct the transcription of the sequence from the gene in the intended tissues of the transformed plant.

For example, for overexpression, a plant promoter fragment may be employed which will direct expression of the gene in all tissues of a regenerated plant. Such promoters are referred to herein as "constitutive" promoters and are active under most environmental conditions and states of development or cell differentiation. Examples of constitutive promoters include the cauliflower mosaic virus (CaMV) 35S

transcription initiation region, the 1'- or 2'- promoter derived from T-DNA of *Agrobacterium tumefaciens*, and other transcription initiation regions from various plant genes known to those of skill.

Alternatively, the plant promoter may direct expression of the
5 polynucleotide of the invention in a specific tissue (tissue-specific promoters) or may be otherwise under more precise environmental control (inducible promoters). Examples of tissue-specific promoters under developmental control include promoters that initiate transcription only in certain tissues, such as fruit, seeds, or flowers. As noted above, the promoters from the *IND1* genes described here are particularly useful for directing gene
10 expression so that a desired gene product is located in the valve margin of fruit. Other suitable promoters include those from genes such as *SHP1* or *SHP2* (Savidge, B., Rounsley, S.D., and Yanofsky, M.F. (1995) *Plant Cell* 7: 721-733). Examples of environmental conditions that may affect transcription by inducible promoters include anaerobic conditions, elevated temperature, or the presence of light.

15 If proper polypeptide expression is desired, a polyadenylation region at the 3'-end of the coding region should be included. The polyadenylation region can be derived from the natural gene, from a variety of other plant genes, or from T-DNA.

The vector comprising the sequences (e.g., promoters or coding regions) from genes of the invention will typically comprise a marker gene that confers a
20 selectable phenotype on plant cells. For example, the marker may encode biocide resistance, particularly antibiotic resistance, such as resistance to kanamycin, G418, bleomycin, hygromycin, or herbicide resistance, such as resistance to chlorosulfuron or Basta.

IND1 nucleic acid sequences of the invention are expressed recombinantly
25 in plant cells to enhance and increase levels of endogenous *IND1* polypeptides. Alternatively, antisense or other *IND1* constructs (described above) are used to suppress *IND1* levels of expression. A variety of different expression constructs, such as expression cassettes and vectors suitable for transformation of plant cells can be prepared. Techniques for transforming a wide variety of higher plant species are well known and
30 described in the technical and scientific literature. See, e.g., Weising *et al. Ann. Rev. Genet.* 22:421-477 (1988). A DNA sequence coding for a *IND1* polypeptide, e.g., a cDNA sequence encoding a full length protein, can be combined with cis-acting (promoter) and trans-acting (enhancer) transcriptional regulatory sequences to direct the

timing, tissue type and levels of transcription in the intended tissues of the transformed plant. Translational control elements can also be used.

The invention provides an *IND1* nucleic acid operably linked to a promoter which, in a preferred embodiment, is capable of driving the transcription of the *IND1* coding sequence in plants. The promoter can be, e.g., derived from plant or viral sources. The promoter can be, e.g., constitutively active, inducible, or tissue specific. In construction of recombinant expression cassettes, vectors, transgenics, of the invention, a different promoters can be chosen and employed to differentially direct gene expression, e.g., in some or all tissues of a plant or animal. Typically, as discussed above, desired promoters are identified by analyzing the 5' sequences of a genomic clone corresponding to the *IND1* genes described here.

A. Constitutive Promoters

A promoter fragment can be employed which will direct expression of *IND1* nucleic acid in all transformed cells or tissues, e.g. as those of a regenerated plant. The term "constitutive regulatory element" means a regulatory element that confers a level of expression upon an operatively linked nucleic molecule that is relatively independent of the cell or tissue type in which the constitutive regulatory element is expressed. A constitutive regulatory element that is expressed in a plant generally is widely expressed in a large number of cell and tissue types. Promoters that drive expression continuously under physiological conditions are referred to as "constitutive" promoters and are active under most environmental conditions and states of development or cell differentiation.

A variety of constitutive regulatory elements useful for ectopic expression in a transgenic plant are well known in the art. The cauliflower mosaic virus 35S (CaMV 35S) promoter, for example, is a well-characterized constitutive regulatory element that produces a high level of expression in all plant tissues (Odell *et al.*, *Nature* 313:810-812 (1985)). The CaMV 35S promoter can be particularly useful due to its activity in numerous diverse plant species (Benfey and Chua, *Science* 250:959-966 (1990); Futterer *et al.*, *Physiol. Plant* 79:154 (1990); Odell *et al.*, *supra*, 1985). A tandem 35S promoter, in which the intrinsic promoter element has been duplicated, confers higher expression levels in comparison to the unmodified 35S promoter (Kay *et al.*, *Science* 236:1299 (1987)). Other useful constitutive regulatory elements include, for example, the cauliflower mosaic virus 19S promoter; the Figwort mosaic virus promoter; and the

nopaline synthase (nos) gene promoter (Singer *et al.*, *Plant Mol. Biol.* 14:433 (1990); An, *Plant Physiol.* 81:86 (1986)).

Additional constitutive regulatory elements including those for efficient expression in monocots also are known in the art, for example, the pEmu promoter and promoters based on the rice Actin-1 5' region (Last *et al.*, *Theor. Appl. Genet.* 81:581 (1991); Mcelroy *et al.*, *Mol. Gen. Genet.* 231:150 (1991); Mcelroy *et al.*, *Plant Cell* 2:163 (1990)). Chimeric regulatory elements, which combine elements from different genes, also can be useful for ectopically expressing a nucleic acid molecule encoding an IND1 polynucleotide (Comai *et al.*, *Plant Mol. Biol.* 15:373 (1990)).

Other examples of constitutive promoters include the 1'- or 2'- promoter derived from T-DNA of *Agrobacterium tumefaciens* (see, e.g., Mengiste (1997) *supra*; O'Grady (1995) *Plant Mol. Biol.* 29:99-108); actin promoters, such as the *Arabidopsis* actin gene promoter (see, e.g., Huang (1997) *Plant Mol. Biol.* 1997 33:125-139); alcohol dehydrogenase (Adh) gene promoters (see, e.g., Millar (1996) *Plant Mol. Biol.* 31:897-904); *ACT11* from *Arabidopsis* (Huang *et al.* *Plant Mol. Biol.* 33:125-139 (1996)), *Cat3* from *Arabidopsis* (GenBank No. U43147, Zhong *et al.*, *Mol. Gen. Genet.* 251:196-203 (1996)), the gene encoding stearyl-acyl carrier protein desaturase from *Brassica napus* (Genbank No. X74782, Solocombe *et al.* *Plant Physiol.* 104:1167-1176 (1994)), *GPc1* from maize (GenBank No. X15596, Martinez *et al.* *J. Mol. Biol.* 208:551-565 (1989)), *Gpc2* from maize (GenBank No. U45855, Manjunath *et al.*, *Plant Mol. Biol.* 33:97-112 (1997)), other transcription initiation regions from various plant genes known to those of skill. See also Holtorf *Plant Mol. Biol.* 29:637-646 (1995).

B. Inducible Promoters

Alternatively, a plant promoter may direct expression of the *IND1* nucleic acid of the invention under the influence of changing environmental conditions or developmental conditions. Examples of environmental conditions that may effect transcription by inducible promoters include anaerobic conditions, elevated temperature, drought, or the presence of light. Such promoters are referred to herein as "inducible" promoters. For example, the invention incorporates the drought-inducible promoter of maize (Busk (1997) *supra*); the cold, drought, and high salt inducible promoter from potato (Kirch (1997) *Plant Mol. Biol.* 33:897-909).

Alternatively, plant promoters which are inducible upon exposure to plant hormones, such as auxins, are used to express the nucleic acids of the invention. For

example, the invention can use the auxin-response elements E1 promoter fragment (AuxREs) in the soybean (*Glycine max* L.) (Liu (1997) *Plant Physiol.* 115:397-407); the auxin-responsive Arabidopsis GST6 promoter (also responsive to salicylic acid and hydrogen peroxide) (Chen (1996) *Plant J.* 10: 955-966); the auxin-inducible parC promoter from tobacco (Sakai (1996) 37:906-913); a plant biotin response element (Streit (1997) *Mol. Plant Microbe Interact.* 10:933-937); and, the promoter responsive to the stress hormone abscisic acid (Sheen (1996) *Science* 274:1900-1902).

Plant promoters which are inducible upon exposure to chemicals reagents which can be applied to the plant, such as herbicides or antibiotics, are also used to express the nucleic acids of the invention. For example, the maize In2-2 promoter, activated by benzenesulfonamide herbicide safeners, can be used (De Veylder (1997) *Plant Cell Physiol.* 38:568-577); application of different herbicide safeners induces distinct gene expression patterns, including expression in the root, hydathodes, and the shoot apical meristem. *IND1* coding sequence can also be under the control of, e.g., a tetracycline-inducible promoter, e.g., as described with transgenic tobacco plants containing the *Avena sativa* L. (oat) arginine decarboxylase gene (Masgrau (1997) *Plant J.* 11:465-473); or, a salicylic acid-responsive element (Stange (1997) *Plant J.* 11:1315-1324; Uknes *et al.*, *Plant Cell* 5:159-169 (1993); Bi *et al.*, *Plant J.* 8:235-245 (1995)).

Particularly useful inducible regulatory elements include copper-inducible regulatory elements (Mett *et al.*, *Proc. Natl. Acad. Sci. USA* 90:4567-4571 (1993); Furst *et al.*, *Cell* 55:705-717 (1988)); tetracycline and chlor-tetracycline-inducible regulatory elements (Gatz *et al.*, *Plant J.* 2:397-404 (1992); Röder *et al.*, *Mol. Gen. Genet.* 243:32-38 (1994); Gatz, *Meth. Cell Biol.* 50:411-424 (1995)); ecdysone inducible regulatory elements (Christopherson *et al.*, *Proc. Natl. Acad. Sci. USA* 89:6314-6318 (1992); Kreutzweiser *et al.*, *Ecotoxicol. Environ. Safety* 28:14-24 (1994)); heat shock inducible regulatory elements (Takahashi *et al.*, *Plant Physiol.* 99:383-390 (1992); Yabe *et al.*, *Plant Cell Physiol.* 35:1207-1219 (1994); Ueda *et al.*, *Mol. Gen. Genet.* 250:533-539 (1996)); and *lac* operon elements, which are used in combination with a constitutively expressed *lac* repressor to confer, for example, IPTG-inducible expression (Wilde *et al.*, *EMBO J.* 11:1251-1259 (1992)). An inducible regulatory element useful in the transgenic plants of the invention also can be, for example, a nitrate-inducible promoter derived from the spinach nitrite reductase gene (Back *et al.*, *Plant Mol. Biol.* 17:9 (1991)) or a light-inducible promoter, such as that associated with the small subunit of RuBP

carboxylase or the LHCP gene families (Feinbaum *et al.*, *Mol. Gen. Genet.* 226:449 (1991); Lam and Chua, *Science* 248:471 (1990)).

C. Tissue-Specific Promoters

5 Alternatively, the plant promoter may direct expression of the polynucleotide of the invention in a specific tissue (tissue-specific promoters). Tissue specific promoters are transcriptional control elements that are only active in particular cells or tissues at specific times during plant development, such as in vegetative tissues or reproductive tissues. Promoters from the *IND1* genes of the invention are particularly
10 useful for tissue-specific direction of gene expression so that a desired gene product is generated only or preferentially in embryos or seeds, as described below.

Examples of tissue-specific promoters under developmental control include promoters that initiate transcription only (or primarily only) in certain tissues, such as vegetative tissues, e.g., roots or leaves, or reproductive tissues, such as fruit,
15 ovules, seeds, pollen, pistils, flowers, or any embryonic tissue. Reproductive tissue-specific promoters may be, e.g., ovule-specific, embryo-specific, endosperm-specific, integument-specific, seed and seed coat-specific, pollen-specific, petal-specific, sepal-specific, or some combination thereof.

The invention provides a transgenic plant that is characterized by delayed
20 seed dispersal due to expression of a nucleic acid molecule encoding an *IND1* gene product, or an antisense construct thereof, operatively linked to a dehiscence zone-selective regulatory element. The dehiscence zone-selective regulatory element can be, for example, an *SHP1* regulatory element or *SHP2* regulatory element. The *SHP1* regulatory element can be derived from the Arabidopsis *SHP1* genomic sequence
25 disclosed herein as SEQ ID NO:5 and can be, for example, a 5' regulatory sequence or intronic regulatory element. Similarly, the *SHP2* regulatory element can be derived from the Arabidopsis *SHP2* genomic sequence disclosed herein as SEQ ID NO:6 and can be, for example, a 5' regulatory sequence or intronic regulatory element.

A dehiscence zone-selective regulatory element can be derived from a
30 gene that is an ortholog of Arabidopsis *IND1* and is selectively expressed in the valve margin or dehiscence zone of a seed plant. A dehiscence zone-selective regulatory element can be derived, for example, from an *IND1* ortholog of the *Brassicaceae*, such as a *Brassica napus*, *Brassica oleracea*, *Brassica campestris*, *Brassica juncea*, *Brassica nigra* or *Brassica carinata* *IND1* ortholog. A dehiscence zone-selective regulatory

element can be derived, for example, from an *IND1* canola ortholog. A dehiscence zone-selective regulatory element also can be derived, for example, from a leguminous *IND1* ortholog, such as a soybean, pea, chickpea, moth bean, broad bean, kidney bean, lima bean, lentil, cowpea, dry bean, peanut, alfalfa, lucerne, birdsfoot trefoil, clover,
5 stylosanthes, *lotononis bainesii*, or sainfoin *IND1* ortholog.

Dehiscence zone-selective regulatory elements also can be derived from a variety of other genes that are selectively expressed in the valve margin or dehiscence zone of a seed plant. For example, the rapeseed gene *RDPG1* is selectively expressed in the dehiscence zone (Petersen *et al.*, *Plant Mol. Biol.* 31:517-527 (1996)). Thus, the
10 *RDPG1* promoter or an active fragment thereof can be a dehiscence zone-selective regulatory element as defined herein. Additional genes such as the rapeseed gene *SAC51* also are known to be selectively expressed in the dehiscence zone; the *SAC51* promoter or an active fragment thereof also can be a dehiscence zone-selective regulatory element of the invention (Coupe *et al.*, *Plant Mol. Biol.* 23:1223-1232 (1993)). The skilled artisan
15 understands that a regulatory element of any such gene selectively expressed in cells of the valve margin or dehiscence zone can be a dehiscence zone-selective regulatory element as defined herein.

Additional dehiscence zone-selective regulatory elements can be identified and isolated using routine methodology. Differential screening strategies using, for
20 example, RNA prepared from the dehiscence zone and RNA prepared from adjacent pod material can be used to isolate cDNAs selectively expressed in cells of the dehiscence zone (Coupe *et al.*, *supra*, 1993); subsequently, the corresponding genes are isolated using the cDNA sequence as a probe.

Enhancer trap or gene trap strategies also can be used to identify and
25 isolate a dehiscence zone-selective regulatory element of the invention (Sundaresan *et al.*, *supra*, 1995; Koncz *et al.*, *Proc. Natl. Acad. Sci. USA* 86:8467-8471 (1989); Kertbundit *et al.*, *Proc. Natl. Acad. Sci. USA* 88:5212-5216 (1991); Topping *et al.*, *Development* 112:1009-1019 (1991)). Enhancer trap elements include a reporter gene such as *GUS* with a weak or minimal promoter, while gene trap elements lack a promoter sequence,
30 relying on transcription from a flanking chromosomal gene for reporter gene expression. Transposable elements included in the constructs mediate fusions to endogenous loci; constructs selectively expressed in the valve margin or dehiscence zone are identified by their pattern of expression. With the inserted element as a tag, the flanking dehiscence zone-selective regulatory element is cloned using, for example, inverse polymerase chain

reaction methodology (see, for example, Aarts *et al.*, *Nature* 363:715-717 (1993); see also, Ochman *et al.*, "Amplification of Flanking Sequences by Inverse PCR," in Innis *et al.*, *supra*, 1990). The Ac/Ds transposition system of Sundaresan *et al.*, *Genes. Devel.* 9:1797-1810 (1995), can be particularly useful in identifying and isolating a dehiscence zone-selective regulatory element of the invention.

Dehiscence zone-selective regulatory elements also can be isolated by inserting a library of random genomic DNA fragments in front of a promoterless reporter gene and screening transgenic plants transformed with the library for dehiscence zone-selective reporter gene expression. The promoterless vector pROA97, which contains the npt gene and the GUS gene each under the control of the minimal 35S promoter, can be useful for such screening. The genomic library can be, for example, Sau3A fragments of *Arabidopsis thaliana* genomic DNA or genomic DNA from, for example, another Brassicaceae of interest (Ott *et al.*, *Mol. Gen. Genet.* 223:169-179 (1990); Claes *et al.*, *The Plant Journal* 1:15-26 (1991)).

Dehiscence zone-selective expression of a regulatory element of the invention can be demonstrated or confirmed by routine techniques, for example, using a reporter gene and in situ expression analysis. The GUS and firefly luciferase reporters are particularly useful for in situ localization of plant gene expression (Jefferson *et al.*, *EMBO J.* 6:3901 (1987); Ow *et al.*, *Science* 334:856 (1986)), and promoterless vectors containing the GUS expression cassette are commercially available, for example, from Clontech (Palo Alto, CA). To identify a dehiscence zone-selective regulatory element of interest such as an *IND1* regulatory element, one or more nucleotide portions of the *IND1* gene can be generated using enzymatic or PCR-based methodology (Glick and Thompson, *supra*, 1993; Innis *et al.*, *supra*, 1990); the resulting segments are fused to a reporter gene such as GUS and analyzed as described above.

Other tissue-specific promoters include seed promoters. Suitable seed-specific promoters are derived from the following genes: *MAC1* from maize (Sheridan (1996) *Genetics* 142:1009-1020); *Cat3* from maize (GenBank No. L05934, Abler (1993) *Plant Mol. Biol.* 22:10131-1038); *viviparous-1* from *Arabidopsis* (Genbank No. U93215); *atmyc1* from *Arabidopsis* (Urao (1996) *Plant Mol. Biol.* 32:571-57; Conceicao (1994) *Plant* 5:493-505); *napA* from *Brassica napus* (GenBank No. J02798, Josefsson (1987) *JBL* 26:12196-1301); and the napin gene family from *Brassica napus* (Sjodahl (1995) *Planta* 197:264-271).

A variety of promoters specifically active in vegetative tissues, such as leaves, stems, roots and tubers, can also be used to express the *IND1* nucleic acids of the invention. For example, promoters controlling patatin, the major storage protein of the potato tuber, can be used, see, e.g., Kim (1994) *Plant Mol. Biol.* 26:603-615; Martin
5 (1997) *Plant J.* 11:53-62. The ORF13 promoter from *Agrobacterium rhizogenes* which exhibits high activity in roots can also be used (Hansen (1997) *Mol. Gen. Genet.* 254:337-343. Other useful vegetative tissue-specific promoters include: the tarin promoter of the gene encoding a globulin from a major taro (*Colocasia esculenta* L. Schott) corm protein family, tarin (Bezerra (1995) *Plant Mol. Biol.* 28:137-144); the
10 curculin promoter active during taro corm development (de Castro (1992) *Plant Cell* 4:1549-1559) and the promoter for the tobacco root-specific gene TobRB7, whose expression is localized to root meristem and immature central cylinder regions (Yamamoto (1991) *Plant Cell* 3:371-382).

Leaf-specific promoters, such as the ribulose biphosphate carboxylase
15 (RBCS) promoters can be used. For example, the tomato RBCS1, RBCS2 and RBCS3A genes are expressed in leaves and light-grown seedlings, only RBCS1 and RBCS2 are expressed in developing tomato fruits (Meier (1997) *FEBS Lett.* 415:91-95). A ribulose biphosphate carboxylase promoters expressed almost exclusively in mesophyll cells in leaf blades and leaf sheaths at high levels, described by Matsuoka (1994) *Plant J.*
20 6:311-319, can be used. Another leaf-specific promoter is the light harvesting chlorophyll a/b binding protein gene promoter, see, e.g., Shiina (1997) *Plant Physiol.* 115:477-483; Casal (1998) *Plant Physiol.* 116:1533-1538. The *Arabidopsis thaliana* myb-related gene promoter (Atmyb5) described by Li (1996) *FEBS Lett.* 379:117-121, is leaf-specific. The Atmyb5 promoter is expressed in developing leaf trichomes, stipules,
25 and epidermal cells on the margins of young rosette and cauline leaves, and in immature seeds. Atmyb5 mRNA appears between fertilization and the 16 cell stage of embryo development and persists beyond the heart stage. A leaf promoter identified in maize by Busk (1997) *Plant J.* 11:1285-1295, can also be used.

Another class of useful vegetative tissue-specific promoters are
30 meristematic (root tip and shoot apex) promoters. For example, the "SHOOTMERISTEMLESS" and "SCARECROW" promoters, which are active in the developing shoot or root apical meristems, described by Di Laurenzio (1996) *Cell* 86:423-433; and, Long (1996) *Nature* 379:66-69; can be used. Another useful promoter is that which controls the expression of 3-hydroxy-3- methylglutaryl coenzyme A

reductase HMG2 gene, whose expression is restricted to meristematic and floral (secretory zone of the stigma, mature pollen grains, gynoecium vascular tissue, and fertilized ovules) tissues (see, e.g., Enjuto (1995) *Plant Cell* 7:517-527). Also useful are kn1-related genes from maize and other species which show meristem-specific expression, see, e.g., Granger (1996) *Plant Mol. Biol.* 31:373-378; Kerstetter (1994) *Plant Cell* 6:1877-1887; Hake (1995) *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* 350:45-51. For example, the *Arabidopsis thaliana* KNAT1 promoter. In the shoot apex, KNAT1 transcript is localized primarily to the shoot apical meristem; the expression of KNAT1 in the shoot meristem decreases during the floral transition and is restricted to the cortex of the inflorescence stem (see, e.g., Lincoln (1994) *Plant Cell* 6:1859-1876).

One of skill will recognize that a tissue-specific promoter may drive expression of operably linked sequences in tissues other than the target tissue. Thus, as used herein a tissue-specific promoter is one that drives expression preferentially in the target tissue, but may also lead to some expression in other tissues as well.

In another embodiment, a *IND1* nucleic acid is expressed through a transposable element. This allows for constitutive, yet periodic and infrequent expression of the constitutively active polypeptide. The invention also provides for use of tissue-specific promoters derived from viruses which can include, e.g., the tobamovirus subgenomic promoter (Kumagai (1995) *Proc. Natl. Acad. Sci. USA* 92:1679-1683; the rice tungro bacilliform virus (RTBV), which replicates only in phloem cells in infected rice plants, with its promoter which drives strong phloem-specific reporter gene expression; the cassava vein mosaic virus (CVMV) promoter, with highest activity in vascular elements, in leaf mesophyll cells, and in root tips (Verdaguer (1996) *Plant Mol. Biol.* 31:1129-1139).

V. Production of transgenic plants

DNA constructs of the invention may be introduced into the genome of the desired plant host by a variety of conventional techniques. For example, the DNA construct may be introduced directly into the genomic DNA of the plant cell using techniques such as electroporation and microinjection of plant cell protoplasts, or the DNA constructs can be introduced directly to plant tissue using ballistic methods, such as DNA particle bombardment. Alternatively, the DNA constructs may be combined with suitable T-DNA flanking regions and introduced into a conventional *Agrobacterium tumefaciens* host vector. The virulence functions of the *Agrobacterium tumefaciens* host

will direct the insertion of the construct and adjacent marker into the plant cell DNA when the cell is infected by the bacteria.

Microinjection techniques are known in the art and well described in the scientific and patent literature. The introduction of DNA constructs using polyethylene glycol precipitation is described in Paszkowski et al. *Embo J.* 3:2717-2722 (1984).
5 Electroporation techniques are described in Fromm et al. *Proc. Natl. Acad. Sci. USA* 82:5824 (1985). Ballistic transformation techniques are described in Klein et al. *Nature* 327:70-73 (1987).

Agrobacterium tumefaciens-mediated transformation techniques, including
10 disarming and use of binary vectors, are well described in the scientific literature. See, for example Horsch et al. *Science* 233:496-498 (1984), and Fraley et al. *Proc. Natl. Acad. Sci. USA* 80:4803 (1983).

Transformed plant cells which are derived by any of the above transformation techniques can be cultured to regenerate a whole plant which possesses the
15 transformed genotype and thus the desired phenotype such as seedlessness. Such regeneration techniques rely on manipulation of certain phytohormones in a tissue culture growth medium, typically relying on a biocide and/or herbicide marker which has been introduced together with the desired nucleotide sequences. Plant regeneration from cultured protoplasts is described in Evans et al., *Protoplasts Isolation and Culture*,
20 *Handbook of Plant Cell Culture*, pp. 124-176, MacMillan Publishing Company, New York, 1983; and Binding, *Regeneration of Plants, Plant Protoplasts*, pp. 21-73, CRC Press, Boca Raton, 1985. Regeneration can also be obtained from plant callus, explants, organs, or parts thereof. Such regeneration techniques are described generally in Klee et al. *Ann. Rev. of Plant Phys.* 38:467-486 (1987).

25 The nucleic acids of the invention can be used to confer desired traits on essentially any plant. Thus, the invention has use over a broad range of plants, including species from the genera Asparagus, Atropa, Avena, Brassica, Citrus, Citrullus, Capsicum, Cucumis, Cucurbita, Daucus, Fragaria, Glycine, Gossypium, Helianthus, Heterocallis, Hordeum, Hyoscyamus, Lactuca, Linum, Lolium, Lycopersicon, Malus, Manihot,
30 Majorana, Medicago, Nicotiana, Oryza, Panieum, Pannasetum, Persea, Pisum, Pyrus, Prunus, Raphanus, Secale, Senecio, Sinapis, Solanum, Sorghum, Trigonella, Triticum, Vitis, Vigna, and, Zea. A useful plant of the invention can be a dehiscent seed plant, and a particularly useful plant of the invention can be a member of the *Brassicaceae*, such as rapeseed, or a member of the *Fabaceae*, such as a soybean, pea, lentil or bean plant.

In one embodiment, the invention provides a dehiscent seed plant that is characterized by delayed seed dispersal due to suppressed expression of a nucleic acid molecule encoding an IND1 gene product in the dehiscent seed plant. As used herein, the term "dehiscent seed plant" means a plant that produces a dry dehiscent fruit, which has fruit walls that open to permit escape of the seeds contained therein. Dehiscent fruits commonly contain several seeds and include the fruits known, for example, as legumes, capsules and siliques.

In one embodiment, the invention provides a plant that is characterized by delayed seed dispersal due to suppressed expression of a nucleic acid molecule encoding an IND1 gene product, where the plant is a member of the *Brassicaceae*. The *Brassicaceae*, commonly known as the Brassicas, are a diverse group of crop plants with great economic value worldwide (see, for example, Williams and Hill, *Science* 232:1385-1389 (1986), which is incorporated herein by reference). The *Brassicaceae* produce seed oils for margarine, salad oil, cooking oil, plastic and industrial uses; condiment mustard; leafy, stored, processed and pickled vegetables; animal fodders and green manures for soil rejuvenation. A particularly useful non-naturally occurring Brassica plant of the invention is the oilseed plant canola.

There are six major Brassica species of economic importance, each containing a range of plant forms. *Brassica napus* includes plants such as the oilseed rapes and rutabaga. *Brassica oleracea* are the cole crops such as cabbage, cauliflower, kale, kohlrabi and Brussels sprouts. *Brassica campestris* (*Brassica rapa*) includes plants such as Chinese cabbage, turnip and pak choi. *Brassica juncea* includes a variety of mustards; *Brassica nigra* is the black mustard; and *Brassica carinata* is Ethiopian mustard. The skilled artisan understands that any member of the *Brassicaceae* can be modified as disclosed herein to produce a non-naturally occurring Brassica plant characterized by delayed seed dispersal.

In a second embodiment, the invention provides a plant that is characterized by delayed seed dispersal due to suppressed expression of a nucleic acid molecule encoding an IND1 gene product, where the plant is a member of the *Fabaceae*. The *Fabaceae*, which are commonly known as members of the pea family, are plants that produce a characteristic dry dehiscent fruit known as a legume. The legume is derived from a single carpel and dehisces along the suture of the carpel margins and along the median vein. The *Fabaceae* encompass both grain legumes and forage legumes. Grain legumes include, for example, soybean (glycine), pea, chickpea, moth bean, broad bean,

kidney bean, lima bean, lentil, cowpea, dry bean and peanut. Forage legumes include alfalfa, lucerne, birdsfoot trefoil, clover, stylosanthes species, *lotononis bainessii* and sainfoin. The skilled artisan will recognize that any member of the *Fabaceae* can be modified as disclosed herein to produce a non-naturally occurring plant of the invention
5 characterized by delayed seed dispersal.

A non-naturally occurring plant of the invention characterized by delayed seed dispersal also can be a member of the plant genus *Cuphea* (family *Lythraceae*). A *Cuphea* plant is particularly valuable since *Cuphea* oilseeds contain industrially and nutritionally important medium-chain fatty acids, especially lauric acid, which is
10 currently supplied only by coconut and palm kernel oils.

A non-naturally occurring plant of the invention also can be, for example, one of the monocotyledonous grasses, which produce many of the valuable small-grain cereal crops of the world. Suppression of *IND1* expression as described above, can be useful in generating a non-naturally occurring small grain cereal plant, such as a barley,
15 wheat, oat, rye, orchard grass, guinea grass, sorghum or turf grass plant characterized by delayed seed dispersal.

VI. Additional modifications that modulate seed dispersal

It should be recognized that a plant of the invention, which contains an
20 exogenous *IND1* polynucleotide, also can contain one or more additional modifications, including naturally and non-naturally occurring modifications, that can modulate the delay in seed dispersal. For example, the plant hormone ethylene promotes fruit dehiscence, and modified expression or activity of positive or negative regulators of the ethylene response can be included in a plant of the invention (see, generally, Meakin and
25 Roberts, *J. Exp. Botany* 41:1003-1011 (1990); Ecker, *Science* 268:667-675 (1995); Chao *et al.*, *Cell* 89:1133-1144 (1997)).

Mutations in positive regulators of the ethylene response show a reduction or absence of responsiveness to treatment with exogenous ethylene. Arabidopsis mutations in positive regulators of the ethylene response include mutations in *etr*, which
30 inactivate a histidine kinase ethylene receptor (Bleeker *et al.*, *Science* 241:1086-1089 (1988); Schaller and Bleeker, *Science* 270:1809-1811 (1995)); *ers* (Hua *et al.*, *Science* 269:1712-1714 (1995)); *ein2* (Guzman and Ecker, *Plant Cell* 2:513 (1990)); *ein3* (Rothenberg and Ecker, *Sem. Dev. Biol. Plant Dev. Genet.* 4:3-13 (1993); Kieber and Ecker, *Trends Genet.* 9:356-362 (1993)); *ain1* (van der Straeten *et al.*, *Plant Physiol.*

102:401-408 (1993)); *eti* (Harpham *et al.*, *An. Bot.* 68:55 (1991)) and *ein4*, *ein5*, *ein6*, and *ein7* (Roman *et al.*, *Genetics* 139: 1393-1409 (1995)). Similar genetic functions are found in other plant species; for example, the never-ripe mutation corresponds to *etr* and confers ethylene insensitivity in tomato (Lanahan *et al.*, *The Plant Cell* 6:521-530 (1994);
5 Wilkinson *et al.*, *Science* 270:1807-1809 (1995)). A plant of the invention can include a modification that results in altered expression or activity of any such positive regulator of the ethylene response. A mutation in a positive regulator, for example, can be included in a plant of the invention and can modify the delay in seed dispersal in such plants, for example, by further postponing the delay in seed dispersal.

10 Mutations in negative regulators of the ethylene response display ethylene responsiveness in the absence of exogenous ethylene. Such mutations include those relating to ethylene overproduction, for example, the *eto1*, *eto2*, and *eto3* mutants, and those relating to constitutive activation of the ethylene signalling pathway, for example, mutations in *CTR1*, a negative regulator with sequence similarity to the Raf family of
15 protein kinases (Kieber *et al.*, *Cell* 72:427-441 (1993), which is incorporated herein by reference). A plant of the invention can include a modification that results in altered expression or activity of any such negative regulator of the ethylene response. A mutation resulting in ethylene responsiveness in the absence of exogenous ethylene, for example, can be included in a non-naturally occurring plant of the invention and can
20 modify, for example, diminish, the delay in seed dispersal.

Fruit morphological mutations also can be included in a plant of the invention. Such mutations include those in carpel identity genes such as *AGAMOUS* (Bowman *et al.*, *supra*, 1989; Yanofsky *et al.*, *supra*, 1990) and in genes required for normal fruit development such as *ETTIN*, *CRABS CLAW*, *SPATULA*, *AGL8* and
25 *TOUSLED* (Sessions *et al.*, *Development* 121:1519-1532 (1995); Alvarez and Smyth, *Flowering Newsletter* 23:12-17 (1997); and Roe *et al.*, *Cell* 75:939-950 (1993)). Thus, it is understood that a plant of the invention can include one or more additional genetic modifications, which can diminish or enhance the delay in seed dispersal.

30 VII. Expression of cytotoxic gene products

The present invention also provides a recombinant nucleic acid molecule that includes a dehiscence zone-selective regulatory element operatively linked to a nucleic acid molecule encoding a cytotoxic gene product. Further provided herein is a plant of the invention that is characterized by delayed seed dispersal due to expression of

a recombinant nucleic acid molecule having a dehiscence zone--selective regulatory element operatively linked to a nucleic acid molecule encoding a cytotoxic gene product.

A cytotoxic gene product is a gene product that causes the death of the cell in which it is expressed and, preferably, does not result in the death of cells other than the cell in which it is expressed. Thus, expression of a cytotoxic gene product from a dehiscence zone-selective regulatory element can be used to ablate the dehiscence zone without disturbing neighboring cells of the replum or valve. A variety of cytotoxic gene products useful in seed plants are known in the art including, for example, diphtheria toxin A chain polypeptides; RNase T1; Barnase RNase; ricin toxin A chain polypeptides; and herpes simplex virus thymidine kinase (tk) gene products. While the diphtheria toxin A chain, RNase T1 and Barnase RNase are preferred cytotoxic gene products, the skilled person recognizes that these, or other cytotoxic gene products can be used with a dehiscence zone-selective regulatory element to generate a non-naturally occurring seed plant characterized by delayed seed dispersal.

Diphtheria toxin is the naturally occurring toxin of *Cornebacterium diphtheriae*, which catalyzes the ADP-ribosylation of elongation factor 2, resulting in inhibition of protein synthesis and consequent cell death (Collier, *Bacteriol. Rev.* 39:54-85 (1975)). A single molecule of the fully active toxin is sufficient to kill a cell (Yamaizumi *et al.*, *Cell* 15:245-250 (1978)). Diphtheria toxin has two subunits: the diphtheria toxin B chain directs internalization to most eukaryotic cells through a specific membrane receptor, whereas the A chain encodes the toxic catalytic domain. The catalytic DT-A chain does not include a signal peptide and is not secreted. Further, any DT-A released from dead cells in the absence of the diphtheria toxin B chain is precluded from cell attachment. Thus, DT-A is cell autonomous and directs killing only of the cells in which it is expressed without apparent damage to neighboring cells. The DT-A expression cassette of Palmiter *et al.*, which contains the 193 residues of the A chain engineered with a synthetic ATG and lacking the native leader sequence, is particularly useful in the seed plants of the invention (Palmiter *et al.*, *Cell* 50:435-443 (1987); Greenfield *et al.*, *Proc. Natl. Acad. Sci., USA* 80:6853-6857 (1983)).

RNase T1 of *Aspergillus oryzae* and Barnase RNase of *Bacillus amylolique-faciens* also are cytotoxic gene products useful in the seed plants of the invention (Thorsness and Nasrallah, *Methods in Cell Biology* 50:439-448 (1995)). Barnase RNase may be more generally toxic to plants than RNase T1 and, thus, is preferred in the methods of the invention.

Ricin, a ribosome-inactivating protein produced by castor bean seeds, also is a cytotoxic gene product useful in a non-naturally occurring seed plant of the invention. The ricin toxin A chain polypeptide can be used to direct cell-specific ablation as described, for example, in Moffat *et al.*, *Development* 114:681-687 (1992). Plant
5 ribosomes are variably susceptible to the plant-derived ricin toxin. The skilled person understands that the toxicity of ricin depends is variable and should be assessed for toxicity in the seed plant species of interest (see Olsnes and Pihl, *Molecular Action of Toxins and Viruses*, pages 51-105, Amsterdam: Elsevier Biomedical Press (1982)).

10

EXAMPLES

The following examples are offered to illustrate, but no to limit the claimed invention.

Example 1:

15

The GT140 valve margin marker (Sundaresan, V., *et al. Genes Dev.* 9, 1797-1810 (1995)) is expressed at the valve margin of the developing gynoecium just prior to fertilization (stage 13) and this pattern persists in the mature fruit (stage 17). As expression of this marker is largely absent from the valve margins of *shp1 shp2* indehiscent fruits (Liljegren, S.J., Ditta, G.S., Eshed, Y., Savidge, B., Bowman, J.L., and
20 Yanofsky, M.F. *Nature*, in press), it was expected that the gene corresponding to this marker might also be involved in valve margin development and be required for fruit dehiscence.

25

To isolate flanking genomic sequence from the GT140 marker insertion site, TAIL/PCR was performed as previously described (Tsugeki, R., *et al. Plant J.* 10, 479-489 (1996)). Subsequent sequencing of the isolated PCR products demonstrated that they correspond to a fully sequenced BAC from chromosome 4, available in the public database as part of the Arabidopsis Genome Initiative. The GT140 insertion is located between two genes, one encoding a predicted basic helix-loop-helix (bHLH) transcription factor and the other representing a novel gene.

30

Through several lines of subsequent investigation, it was confirmed that the bHLH transcription factor (herein referred to as *IND1* as noted below) was the relevant gene (SEQ ID NO:1). Promoter/enhancer::GUS fusions of the *IND1* gene were introduced into wild-type plants and found to express GUS in an identical pattern to that of the GT140 marker line. Interestingly, approximately 25% of the transgenic lines failed

to express significant GUS activity and displayed an indehiscent phenotype. The most likely explanation of these results is that the *IND1::GUS* fusions, as well as of the endogenous *IND1* gene, were cosuppressed. Subsequent RNA blotting confirmed a down regulation of the *IND1* gene in these lines, and further RNA blotting showed, as expected,
5 a decrease in *IND1* gene expression in *shp1 shp2* fruits.

In parallel to the studies of the GT140 valve margin marker described above, screens for Arabidopsis mutants producing indehiscent fruits were also carried out. Besides obtaining additional alleles of *SHP1* and *SHP2* through EMS mutagenesis of *shp2-1* and *shp1-1* seed stocks, indehiscent mutants that were not allelic to either *SHP1* or
10 *SHP2*, respectively were also obtained. Because the GT140 studies suggested the possibility that one or more of these indehiscent mutants might correspond to the *IND1* gene, *IND1* from several of these mutants was cloned and sequenced. Four alleles represent independent mutant alleles of *IND1*. The strongest allele, *ind1-2*, contains a single nucleotide deletion within codon 55 that results in a frameshift and production of a
15 truncated protein of 64 rather than 198 amino acids. The *ind1-1* and *ind1-3* alleles contain nucleotide substitutions at codons 141 and 128 that changes a leucine amino acid to a phenylalanine and an arginine to a histidine, respectively. These affected amino acids are both at conserved positions within the bHLH domain. The *ind1-4* allele contains a
20 nucleotide substitution at codon 92 that changes a glutamine to a stop codon, causing production of a truncated protein of 91 amino acids. Since inactivation of this bHLH transcription factor prevents fruit dehiscence, the gene is referred to as *INDEHISCENT1* (*IND1*) and the mutant as, *ind1*. To date, *ind1* represents the only reported single gene mutation in Arabidopsis that specifically blocks fruit dehiscence.

25 It is understood that the examples and embodiments described herein are for illustrative purposes only and that various modifications or changes in light thereof will be suggested to persons skilled in the art and are to be included within the spirit and purview of this application and scope of the appended claims. All publications, patents, and patent applications cited herein are hereby incorporated by reference in their entirety
30 for all purposes.

WHAT IS CLAIMED IS:

- 1 1. An isolated nucleic acid, comprising an IND1 polynucleotide
2 sequence encoding an IND1 polypeptide at least about 70% identical to SEQ ID NO:2.
- 1 2. The isolated nucleic acid of claim 1, wherein the IND1 polypeptide
2 comprises SEQ ID NO:2.
- 1 3. The isolated nucleic acid of claim 1, wherein the IND1
2 polynucleotide comprises positions from about 2765 to about 3361 of SEQ ID NO 1.
- 1 4. The isolated nucleic acid of claim 3, wherein the IND1
2 polynucleotide comprises SEQ ID NO:1.
- 1 5. An expression cassette comprising a promoter operably linked to
2 an IND1 polynucleotide sequence, or a complement thereof, encoding an IND1
3 polypeptide at least about 70% identical to SEQ ID NO:1.
- 1 6. The expression cassette of claim 5, wherein the IND1 polypeptide
2 comprises SEQ ID NO:2.
- 1 7. The expression cassette of claim 5, wherein the IND1
2 polynucleotide comprises positions from about 2765 to about 3361 of SEQ ID NO 1.
- 1 8. The expression cassette of claim 7, wherein the IND1
2 polynucleotide comprises SEQ ID NO 1.
- 1 9. The expression cassette of claim 5, wherein the promoter is
2 constitutive.
- 1 10. The expression cassette of claim 5, wherein the promoter is tissue
2 specific.
- 1 11. The expression cassette of claim 10, wherein the promoter is a
2 dehiscence zone specific promoter.
- 1 12. The expression cassette of claim 11, wherein the promoter
2 comprises positions from about 1 to about 2764 or from about 3362 to about 3856 of SEQ
3 ID NO:1.

1 13. A plant comprising a recombinant expression cassette comprising a
2 promoter operably linked to a polynucleotide sequence encoding an IND1 polypeptide at
3 least about 70% identical to SEQ ID NO: 1.

1 14. The plant of claim 13, wherein the polynucleotide sequence
2 encoding the IND1 polypeptide is operably linked to the promoter in the antisense
3 orientation.

1 15. The plant of claim 13, wherein the polynucleotide sequence
2 encoding the IND1 polypeptide is operably linked to the promoter in the sense
3 orientation.

1 16. The plant of claim 15, wherein the polynucleotide sequence further
2 comprises a second polynucleotide sequence encoding the IND1 polypeptide wherein the
3 second polynucleotide sequence is operably linked to a second promoter in the antisense
4 orientation.

1 17. The plant of claim 13, wherein lignification is reduced in valve
2 margin cells.

1 18. The plant of claim 13, wherein the promoter is a dehiscence zone-
2 selective regulatory element.

 19. The plant of claim 18, where in the regulatory element comprises
 positions from about 1 to about 2764 or from about 3362 to about 3856 of SEQ ID NO:1.

1 20. A method of delaying fruit dehiscence in a plant, the method
2 comprising suppressing expression of an IND1 nucleic acid in the plant by introducing
3 into the plant a recombinant expression cassette comprising a promoter operably linked to
4 a polynucleotide sequence encoding an IND1 polypeptide at least about 70% identical to
5 SEQ ID NO: 2.


1 21. The method of claim 20, wherein the IND1 polypeptide comprises
2 SEQ ID NO:2.

1 22. The method of claim 20, wherein the IND1 polynucleotide
2 comprises positions from about 2765 to about 3361 of SEQ ID NO:1.

- 1 23. The method of claim 20, wherein the IND1 polynucleotide
2 comprises SEQ ID NO:1.
- 1 24. The method of claim 20, wherein the polynucleotide sequence
2 encoding the IND1 polypeptide is operably linked to the promoter in the antisense
3 orientation.
- 1 25. The method of claim 20, wherein the polynucleotide sequence
2 encoding the IND1 polypeptide is operably linked to the promoter in the sense
3 orientation.
- 1 26. The method of claim 25, wherein the polynucleotide sequence
2 further comprises a second polynucleotide sequence encoding the IND1 polypeptide
3 wherein the second polynucleotide sequence is operably linked to a second promoter in
4 the antisense orientation.
- 1 27. The method of claim 20, wherein lignification is reduced in valve
2 margin cells.
- 1 28. The method of claim 20, wherein the promoter is a dehiscence
2 zone-selective regulatory element.
- 1 29. The method of claim 28, where in the regulatory element
2 comprises positions from about 1 to about 2764 or from about 3362 to about 3856 of SEQ
3 ID NO:1.
- 1 30. The method of claim 20, wherein the recombinant expression
2 cassette is introduced into the plant using *Agrobacterium*.
- 1 31. A method of delaying fruit dehiscence in a plant, the method
2 comprising suppressing expression of an IND1 gene in the plant by introducing into the
3 plant a recombinant expression cassette comprising a polynucleotide sequence at least
4 about 70% identical to positions from about 1 to about 2764 or from about 3362 to about
5 3856 of SEQ ID NO:1.

1 32. The method of claim 31, wherein the polynucleotide sequence
2 comprises positions from about 1 to 2764 or from about 3362 to about 3856 of SEQ ID
3 NO:1.

1 33. The method of claim 31, wherein lignification is reduced in valve
2 margin cells.



SEQUENCE LISTING

SEQ ID NO:1 IND1 genomic

Sequence Range: 1 to 3856

5 CTCTAGACCA TCTACTATCC GGTGTGTGAC CCTTAAAGCT TTTGAAGACT ACTAGAATAA 60

10 TGCAAATACC ATATGTCCAT ATCCATCCTT TTCTTTTGTT TGAAGTGAAC ATTCTAATTT 120

15 TGTAAGAGAA AAAACCTTAT GTTAATATCA CCGTAGGCAA AAAAAATATC TCATCATATT 180

20 AAATTTTAT TATAAGATTA TACATTCTCT CGTTGTAAGA GTTACTCCAA TTGCAAGTGT 240

25 TGTATTAAT AATAAAAAGG ACGAAAGTAG GAAGCTTATA ATTAATTGAT GTTGCATAGT 300

30 ACTGGTATAT TGTGATGAA TATAACAAGT ATGAACATTA ATGCATGAAA CGGGGTATTT 360

35 TGTCTTGAAC TCATTAAAGG CAATGTGAAA AGAAGATGTG AGGTCTCATT TTGAAAATTT 420

40 ATCTTCTAGC TTTGTGATT TTAATCTAT GAAATGAACG CAACATATAG AAATTTTCATG 480

45 TGGACAACGA CATTTAGACG GTATCTTAAT TAGACCGATT AATTAGTAAT ATACTTATAT 540

50 ATATAATTAG TGGTGATTAT AAGTTTACTT ATCCACTTGA GAAATTAAAC AATGGGCAAT 600

55 ACCTTAATGT CGAAGAAGC CGTCCCACT TCGTGTAATG AGTTATGGGG GAGAGATCCT 660

60 GTTAAATCGT CAAATAAAC AACTTAAGAA CTAGAAATTG ACACCAAAAA TCATAAGAG 720

65 AACGTTGAAG AAGTCATTTA TCGTATCCAG CTCATATTTT CTAGCTAAGA TCAAATCAAG 780

70 GCCGTTGAAA GGGCTTGTA GAAATGTCG AAGAAACCGT GGGGTTTAGA AGAAAGACAA 840

75 GAAATAGAAG AACAATGATG TTAAATTGCC TATTTTGGTG TATAGGAGTT GTCAAAAGAG 900

80 GAGAGAGAGA AGAAAATTAG GTCAAAATAA TGAGCACTAA AAATGGAGAC ATGTGTTGAG 960

85 TAACTATTAC AAGAGCGACT TATGCTTCCT TATGGCAATG ATATCCAAAC CAAAGTGCAA 1020

90 CGCTCCTTTT TTGCCCTAAT TTCGTAAAGT CTCTCTCCTT CTTCGTCCTT AGGAAAAACC 1080

95 CTAGAAATTT AATCCCTTGT TCTTGATCTT GCTTTTGTAG TAACCATGAT TTTGACCACA 1140

100 CACTATTTCT TCTATCTTTT GTGGTCTATA GGATTTTGCT TTATATGTGT TTCTTGATTT 1200

105 GCTCCGTACG TACGTATACG AATTAAATG GTTATAACAA GGTTTATATA AACTAGCACA 1260

1320
AATGAGTCCA TGAAATTTGT TAGCGAAAAA GGTAGAAATA TATTGAGTCT TTAAACGGCA
5
1380
ATATATATAA TTTTGCTGCA AAAGTTAGCT TTAATCATGA TCTAATGATA TTTTCTTTAA
1440
10 TTTCTTTTGC CAAATTAATC ACATGCACGG ATTTTGGCA AGTTATGTGT CGAATTCCTC
1500
CATTCACACA AACTTAACT TAATTAGAAC TCTAGGAAAT ATTTTAAAT GACAACCTTA
1560
15 TCGAAAAAAA TTTAGTTATG AAAACAATTC CAGAATTAA CATGAGCTAT ATAATTTAAG
1620
ATAAAATGAA GTAATATTGA TATGTATGTA ATAACATATC TGATTGCGGT AAAAAAAGC
1680
20 ATATCTGATT AAATTGTTCA TGCAGGCCCA TGTCATATG ATGTCATCAC GTTTTTATT
1740
25 TCACAATAAC TAATATATAT TCAAAAAAT AGTTTGTCA GATTAAATTT TTTTGGTGG
1800
TCAGCTTTCT CCAACCTACT AAAGTAGTTT GGAATGTTCT CTTCTTTATT TTTCTTTTCT
1860
30 TTGATTTCTT ATGTTTTTTA TTTATGGAAT TTAAGACGG ATTGTTTAGG TCGTTTCTCT
1920
CTTTTCTTGT TTTCTAAAGT TACTTTTGTA AACTCATCTC CTCCCAATTA GACAGTCAAT
1980
35 CATATAGTTA TCTTTTAATA TATGTCTAGT TGATAAAAAA AATGAAAAA TACTGGTGGT
2040
40 AGTTCTACTA ATGTTTGTGT AAAAAATCTG ATATTATGAA TCTAATCAAT TTCTTTGATC
2100
GTATAATGTG GGTAAATTT AGTAATTTT TACATAAATA AGAACTGTAA TGTGATGTA
2160
45 TATTGGGGAA TCAGTATATT AGCTTGGGTA ACTATACTTC TGGAAATACT TGAAGATTTA
2220
ACTATTTGCA AAATTATAAT TTAGTCCCGA AAAATACAGA CGACGGGACA CGACAACATA
2280
50 TAAGCAGGTT TGAATCTTG AAAATTTTGT ATACATAACC TATATAAATA CTAATGTTCT
2340
55 GGTGCGGTT AAAAGCCTT TCAAAAGTTC CATTTTTTAA ATTCAAGGAC ATTTTACATA
2400
GGAAATAAGT TGAGTCATAA AAAATAATGG TTATTTTGTA AGGTTTTTTT TTTGATTAAA
2460
60 ACGCACATAT TAAGAAGTTA GTTTTTTTT ACTACCAAAT ATCAATTAAT TTAACCAT
2520
GCAACCATTC ATAAACAAT ACTATTAAAG AATATAAATA ATCACAATTT ATTAAATACA
2580
65 CTTAAATTT ACATATAAAT TTACAAAACA TCTAATTAAT TGAAACAGAA AGGAAAGGT
2640

5 AAAATATATC ATAAAATGAG ACATATATCC TATAAAAAAA AAATGAGGCA TATGAAGTAA
2700
ATAATAAGAG ACATGCATGT AAGCATTCCG TTAATTAATC GAGTCAAAGA TATATATCAG
2760
TAAATACATA TGTGTATATT TCTGGAAAAA GAATATATAT ATTGAGAAAT AAGAAAAGAT
2820
10 GAAAATGGAA AATGGTATGT ATAAAAAGAA AGGAGTGTGC GACTCTTGTG TCTCGTCCAA
M E N G M Y K K K G V C D S C V S S K>
2880
15 AAGCAGATCC AACCACAGCC CAAAAGAAG CATGATGGAG CCTCAGCCTC ACCATCTCCT
S R S N H S P K R S M M E P Q P H H L L>
2940
20 CATGGATTGG AACAAAGCTA ATGATCTTCT CACACAAGAA CACGCAGCTT TTCTCAATGA
M D W N K A N D L L T Q E H A A F L N D>
3000
TCCTCACCAT CTCATGTTAG ATCCACCTCC CGAAACCCTA ATTCACCTGG ACGAAGACGA
P H H L M L D P P P E T L I H L D E D E>
3060
25 AGAGTACGAT GAAGACATGG ATGCGATGAA GGAGATGCAG TACATGATCG CCGTCATGCA
E Y D E D M D A M K E M Q Y M I A V M Q>
3120
30 GCCCGTAGAC ATCGACCCTG CCACGGTCCC TAAGCCGAAC CGCCGTAACG TAAGGATAAG
P V D I D P A T V P K P N R R N V R I S>
3180
35 CGACGATCCT CAGACGGTGG TTGCTCGTCG GCGTCGGGAA AGGATCAGCG AGAAGATCCG
D...D P Q T V V A R R R R E R I S E K I R>
3240
40 AATTCTCAAG AGGATCGTGC CTGGTGGTGC GAAGATGGAC ACAGCTTCCA TGCTCGACGA
I L K R I V P G G A K M D T A S M L D E>
3300
AGCCATACGT TACACCAAGT TCTTGAAACG GCAGGTGAGG ATTCTTCAGC CTCACTCTCA
A I R Y T K F L K R Q V R I L Q P H S Q>
3360
45 GATTGGAGCT CCTATGGCTA ACCCCTCTTA CCTTTGTTAT TACCACAACCT CCCAACCCCTG
I G A P M A N P S Y L C Y Y H N S Q P *>
3420
50 ATGAACTACA CAGAAGCTCG CTAGCTAGAC ATTTGGTGTC ATCCTCTCAA CCTTTTTCAT
3480
GTTGATATAT TATATATAGA TGCATAAAGA TTCGATCCAA GATTGTATGG GTGTTTTAAT
3540
55 ATTATTATTC TAAGATATAT GATGTACAAT TGTGTACCAA GTTTCTTTAT CTTGATATCA
3600
60 TATGCATAAA TAATTGGTGA ATAAAAAGAA GATATTGATT GTAAACAAAA AAAAGAAGAT
3660
ATTGATTGTT AATTAGGTT TGATCATTCT GTATGAAAGC TTTGGCCTGC AAATTAATTT
3720
65 TCGATATATA TATATATATA TGGAGAATAT ATATCAAATA CTTTTTTAAT TTGACTATAA
3780
TTTGTATCAA TTATCTGAAT CTGATGAGTG TAGGTTATAT ATGGATTAGC AAAAAAGAAA

3840
ACAACCATTA TTACGCACCT ACATTAAAAA TCATCCACCA AAGAAGAAAC CATCCTCAAG
5 AGGGTTCCTT CTAGAG

SEQ ID NO:2 IND1 protein

10 M E N G M Y K K K G V C D S C V S S K
S R S N H S P K R S M M E P Q P H H L L
M D W N K A N D L L T Q E H A A F L N D
15 P H H L M L D P P P E T L I H L D E D E
E Y D E D M D A M K E M Q Y M I A V M Q
20 P V D I D P A T V P K P N R R N V R I S
D D P Q T V V A R R R R E R I S E K I R
I L K R I V P G G A K M D T A S M L D E
25 A I R Y T K F L K R Q V R I L Q P H S Q
I G A P M A N P S Y L C Y Y H N S Q P *

SEQ ID NO:3 IND1 5' promoter

30 60 CTCTAGACCA TCTACTATCC GGTGTGAC CCTTAAAGCT TTTGAAGACT ACTAGAATTA
120
35 TGCAAATACC ATATGTCCAT ATCCATCCTT TTCTTTGTT TGAAGTGAAC ATTCTAATTT
180
TGTAAGAGAA AAAACCTTAT GTTAATATCA CCGTAGGCAA AAAAAATATC TCATCATATT
240
40 AAATTTTAT TATAAGATTA TACATTCTCT CGTTGTAAGA GTTACTCCAA TTGCAAGTGT
300
TGTATTAAT AATAAAAAGG ACGAAAGTAG GAAGCTTATA ATTAATTGAT GTTGCATAGT
360
45 ACTGGTATAT TGTGATGAA TATAACAAGT ATGAACATTA ATGCATGAAA CGGGGTATTT
420
50 TGTCTTGAAC TCATTAAAGG CAATGTGAAA AGAAGATGTG AGGTCTCATT TTGAAAATTT
480
ATCTTCTAGC TTTGTCGATT TTAAATCTAT GAAATGAACG CAACATATAG AAATTCATG
540
55 TGGACAACGA CATTAGACG GTATCTTAAT TAGACCGATT AATTAGTAAT ATACTTATAT
600
ATATAATTAG TGGTGATTAT AAGTTTACTT ATCCACTTGA GAATTTAAAC AATGGGCAAT
660
60 ACCTTAATGT CGAAGAAGC CGTCCCACT TCGTGTAATG AGTTATGGGG GAGAGATCCT
720
GTAAATCGT CAAATAAAAC AACTTAAGAA CTAGAAATTG ACACCAAAAA TCATAAAGAG

780
AACGTTGAAG AAGTCATTTA TCGTATCCAG CTCATATTTT CTAGCTAAGA TCAAATCAAG
5 840
GCCGTTGAAA GGGCTTGTA GAAATGTGCG AAGAAACCGT GGGGTTTAGA AGAAAGACAA
900
GAAATAGAAG AACAAATGATG TTAAATTGCC TATTTTGGTG TATAGGAGTT GTCAAAGAG
10 960
GAGAGAGAGA AGAAAATTAG GTCAAATAA TGAGCACTAA AAATGGAGAC ATGTGTTGAG
1020
TAACTATTAC AAGAGCGACT TATGCTTCCT TATGGCAATG ATATCCAAAC CAAAGTGCAA
15 1080
CGCTCCTTTT TTGCCCTAAT TTCGTAAAGT CTCTCTCCTT CTTCGTCCTT AGGAAAAACC
1140
CTAGAAATTT AATCCCTTGT TCTTGATCTT GCTTTTGGAG TAACCATGAT TTTGACCACA
1200
CACTATTTCT TCTATCTTTT GTGGTCTATA GGATTTTGCT TTATATGTGT TTCTTGATT
25 1260
GCTCCGTACG TACGTATACG AATTTAAATG GTTATAACAA GGTATATATA AACTAGCACA
1320
AATGAGTCCA TGAAATTTGT TAGCGAAAAA GGTAGAAATA TATTGAGTCT TTAACGGCA
1380
ATATATATAA TTTTGCTGCA AAAGTTAGCT TTAATCATGA TCTAATGATA TTTCTTTAA
35 1440
TTTCCTTTGC CAAATTAATC ACATGCACGG ATTTTGGCA AGTTATGTGT CGAATTCCTC
1500
CATTACACA AACTAAACT TAATTAGAAC TCTAGGAAAT ATTTTAAAT GACAACTTA
40 1560
TCGAAAAAAA TTTAGTTATG AAAACAATTC CAGAATTAAA CATGAGCTAT ATAATTTAAG
1620
ATAAATGAA GTAATATTGA TATGTATGTA ATAACATATC TGATTGCGGT AAAAAAAAC
45 1680
ATATCTGATT AAATTGTTCA TGCAGGCCCA TGCACTATG ATGTCATCAC GTTTTATTT
1740
TCACAATAAC TAATATATAT TCAAAAAAAT AGTTTGTCA GATTAAATT TTTTGGTGG
50 1800
TCAGCTTTCT CCAACCTACT AAAGTAGTTT GGAATGTTCT CTCTTTTATT TTTCTTTTC
1860
TTGATTCTT ATGTTTTTTA TTTATGGAAT TTTAAGACGG ATTGTTTAGG TCGTTCTCT
1920
CTTTCTTGT TTTCTAAAGT TACTTTTGTA AACTCATCTC CTCCCAATTA GACAGTCAAT
1980
CATATAGTTA TCTTTTAATA TATGTCTAGT TGATAAAAAA AATGAAAAA TACTGGTGGT
60 2040
AGTTCTACTA ATGTTTGTGT AAAAAATCTG ATATTATGAA TCTAATCAAT TTCTTTGATC
2100

GTATAATGTG GGTAAATTT AGTAATTTT TACATAAATA AGAACTGTAA TGTTGATGTA
 2160
 TATTGGGGAA TCAGTATATT AGCTTGGGTA ACTATACTTC TGGAAATACT TGAAGATTTA
 5 2220
 ACTATTTGCA AAATTATAAT TTAGTCCCGA AAAATACAGA CGACGGGACA CGACAACATA
 2280
 TAAGCAGGTT TGAATCTTGG AAAATTTTGT ATACATAACC TATATAAATA CTAATGTTCT
 10 2340
 GGTGGGTTT AAAAGCCTTT TCAAAAGTTC CATTTTTAA ATTCAAGGAC ATTTTACATA
 2400
 GGAAATAAGT TGAGTCATAA AAAATAATGG TTATTTTGTA AGGTTTTTTT TTTGATTAAA
 15 2460
 ACGCACATAT TAAGAAGTTA GTTTTTTTC ACTACCAAAT ATCAATTAAT TTTAAACCAT
 20 2520
 GCAACCATTG ATAAAACAAT ACTATTAAAG AATATAAATA ATCACAAAAT ATTAAATACA
 2580
 CTTAAATTT ACATATAAAT TTACAAAACA TCTAATTAAT TGAAACAGAA AGGAAAAGGT
 25 2640
 AAAATATATC ATAAATGAG ACATATATCC TATAAAAAAA AAATGAGGCA TATGAAGTAA
 2700
 ATAATAAGAG ACATGCATGT AAGCATTTCG TTAATTAATC GAGTCAAAGA TATATATCAG
 2760
 TAAATACATA TGTGTATATT TCTGGAAAAA GAATATATAT ATTGAGAAAT AAGAAAAGAT
 35 GAAAA

SEQ ID NO:4 IND1 3' promoter

3420
 40 ATGAACTACA CAGAAGCTCG CTAGCTAGAC ATTTGGTGTC ATCCTCTCAA CCTTTTTCAT
 3480
 GTTGATATAT TATATATAGA TGCATAAAGA TTCGATCCAA GATTGTATGG GTGTTTAAAT
 45 3540
 ATTATTATTC TAAGATATAT GATGTACAAT TGTGTACCAA GTTCTTTTAT CTTGATATCA
 3600
 TATGCATAAA TAATTGGTGA ATAAAAGAA GATATTGATT GTAAACAAA AAAAGAAGAT
 50 3660
 ATTGATTGTT AATTAGGGTT TGATCATTCT GTATGAAAGC TTTGGCCTGC AAATTAATT
 3720
 55 TCGATATATA TATATATATA TGGAGAATAT ATATCAAATA CTTTTTAAAT TTGACTATAA
 3780
 TTTGTATCAA TTATCTGAAT CTGATGAGTG TAGGTTATAT ATGGATTAGC AAAAAAGAAA
 3840
 60 ACAACCATTG TTACGCACCT ACATTAAAAA TCATCCACCA AAGAAGAAAC CTCCTCAAG
 AGGGTCCCT CTAGAG

SEQ ID NO:5 SHP1 genomic

60
AGATCTGCAA CAGTGAAAAG AGAAAACAAA ATGGACTTGA AGAGGTTTGG ACAATGCCAG
5 120
AGATAATGCT TATTCCTAA TATGTTGCCA GCCAAGTGTC AAATTGGCTT TTAAATATG
180
GATTTCTGTA TCAGTGGTCA TATTTGTGGA TCCAACGTAT TCATCATCAA GTTCTCAAGT
10 240
TTGCTTTCAG TGCAATTCTA ATTCACACGT TTAACTTTAA CATGCATGTC ATTATAATTA
300
15 CTTCTTCACT AAGACACAAT ACGGCAAACC TTTCAGATTA TATTAATCTC CATAAATGAA
360
ATAATTAACC TCATAATCAA GATTCAATGT TTCTAAATAT ATATGGACAA AATTTACACG
420
20 GAAGATTAGA TACGTATATT AGTAGATTTA GTCTTTCGTT TGTGCGATAA GATTAACCAC
480
25 CTCATAGATA GTAATATCAT TGTCAAATTC CTCTCGGTTT AGTCGCTAAA TTGTATCTTT
540
TTTAAGCCTA AAAGTAGTGT ATTCGCATAT GACTTATCGT CCTAACTTTT TTTTAAATTA
600
30 ACAAAAAAAT CGAAAAGAAA ATAATCTGTT AAATATTTT TAAGTACTCC ATTAAGTTTA
660
GTTTCTATTT AAAAAATGCT TGAAATTGA CAGTTATGTT CAACAATTTT GAATCATGAG
720
35 CGATGTCTAG ATACTCAGAA TTTAATCAAG ATGTCTTATC AAATTTGTTG TCACTCGAGG
780
40 ACCCACGCAA AAGAAAAGAC TAATATGATT TTTATTTGGT CTGGATATTT TTGTAGAGGA
840
TGAAACTAAG AGAGTGAAAG ATTCGAAATC CACAATGTTC AAGAGAGCTC AAAGCAAAAA
900
45 GAAAAATGAA GATGAAGGAC TAAAGAACAA TAAGCAACTA CTTATACCCT ATTTCCATAA
960
AGGATTCAGG TACTAGGAGA AGTTGAGGCA AGTTMNNNN NATTGATTCA AATTTTCATT
1020
50 TATTTTACA ATTTAATTCA CCTAAGTTAT TATGCATTTC TCATCATTGG TACATTTTCT
1080
55 GTATAGCGTA TTACATATA TGAAATAAAT TAAATATGTC CTCACGTTGC AAGTAGTTAA
1140
TGAATGTCCC CACGCAAAAA AAAATCCCTC CAAATATGTC CACCTTTTCT TTTCTTTTAA
1200
60 ATTCCAAAT TACCATAAAC TTTTGGTTTA CAAAAGATT CTAGAAATTG AGGAAGATAT
1260
CCTAAATGAT TCATGAATCC TTCAATAATC TGAAGTTTGC GATATTTTCG ATTTTCTTCA
1320
65 AGAGTTGCGA TATTTGTAAT TTGGTGACCT TAACTTTTT TTGATAAAGA GTAAACGTTT

1380
TTTCTTAAAA GTAAACTTGG ATTTTATGTT TTAGGGTTCT AGCTCAACTT TGTATTATAT
1440
5 TTCTTGCAAA AAGAGTTCGT TAACTGCATT CTTCAACACT ATAAAGTGAT TATCAAAAAC
1500
ATCTTCATGA ACATTAGAA AAACAATATT TGGTTTCGGT TAGAGCTTGG TTTTGCTTGG
1560
10 CTTGATTCAC ATACCCATTC TAGACTTTGG CATAAATTG ATACGATAGA GAGTATCTAA
1620
15 TGGTAATGCA GAAGGGTAAA AAAAGGAAGA GAGAAAAGGT GAGAAAGATT ACCAAAAATA
1680
AGGAGTTTCA AAAGATGGTT CTGATGAGAA ACAGAGCCCA TCCCTCTCCT TTTCCCTTC
1740
20 CCATGAAAGA AATCGGATGG TCCTCCTTCA ATGTCCTCCA CCTACTCTTC TCTCTTTTCT
1800
TTTTTCTTT CTTATTATTA ACCATTTAAT TAATTTCCCC TTCAATTTC GTTTCTAGTT
1860
25 CTGTAAAAAG AAAATACACA TCTCACTTAT AGATATCCAT ATCTATTTAT ATGCATGTAT
1920
30 AGAGAATAAA AAAGTGTGAG TTTCTAGGTA TGTGAGTAT GTGCTGTTG GACAATTGTT
1980
AGATGATCTG TCCATTTTTT TCTTTTTTCT TCTGTGTATA AATATATTG AGCACAAGA
2040
35 AAAACTAATA ACCTTCTGTT TTCAGCAACT AGGGTCTTAT AACCTTCAA GAAATATTC
2100
TTCAATTGAA AACCCATAAA CCAAAATAGA TATTACAAA GGAAAGAGAG ATATTTTCAA
2160
40 GAACAACATA ATTAGAAAAG CAGAAGCAGC AGTTAAGTGG TACTGAGATA AATGATATAG
2220
45 TTTCTCTTCA AGAACAGTTT CTCATTACCC ACCTTCTCCT TTTTGCTGAT CTATCGTAAT
2280
CTTGAGAACT CAGGTAAGGT TGTGAATATT ATGCACCATT CATTAAACCCT AAAAATAAGA
2340
50 GATTTAAAT AAATGTTTCT TCTTTCTCTG ATTCTTGTGT AACCAATTCA TGGGTTTGAT
2400
ATGTTTCTTG GTTATTGCTT ATCAACAAAG AGATTGATC ATTATAAAGT AGATTAATAA
2460
55 CTCTTAAACA CACAAAGTTT CTTTATTTTT TAGTTACATC CCTAATTCTA GACCAGAACA
2520
60 TGGATTGAT CTATTTCTTG GTTATGTATC TTGATCAGGA AAAGGGATTT GATCATCAAG
2580
ATTAGCCTTC TCTCTCTCTC TCTAGATATC TTTCTTGAAT TTAGAAATCT TTATTTAATT
2640
65 ATTTGGTGAT GTCATATATG GATCAATGGA GGAAGGTGGG AGTAGTCAGC ACGCAGAGAG
2700
TAGCAAGAAA CTAGGGAGAG GGAAAATAGA GATAAGAGG ATAGAGAACA CAACAAATCG

2760
TCAAGTTACT TTCTGCAAAC GACGCAATGG TCTTCTCAAG AAAGCTTATG AACTCTCTGT
5 CTTGTGTGAT GCCGAAGTTG CCCTCGTCAT CTTCTCCACT CGTGGCCGTC TCTATGAGTA
2820
CGCCAACAAC AGGTACGCTT CTCCTACTCT ATTTCTTGAT CTTGTTTTCT TAATTTTAAC
10 TAAACAAGAT CCTAGTTCAA ATGATAACAA AGTGGGGATT GAGAGCCAAG ATTAGGGTTT
2940
GGTTAATTTA GAAAACCAGA TTCTACTTGT TGATACATTT AATATCTCTC TAGCTAGATT
15 TAGTACTCTC TCCTCTATAT ATGTGTGGGT GTGTGTGTAA GTGTGTATAT GTATGCAAAT
3060
GCAAGAAGAA GAAGAAAAAG TTATCTTGTC TTCTCAAATT CTGATCAGCT TTGACCTTAG
20 TTTCACTCTT TTTTCTGCAA ATCATTTGAA CCTGATGCAT GTCAGTTTCT ACAATACACT
3180
TTTAATTTTG ACGGCCCATC AAATTTCTTA GGGTTTACTT CAGTGAACAA AATTGGGTTC
25 TTGACACGAT TTAGCATGTA TATATAAAAA TAGGGGATGA TCAAGACTTA TGTAACCTCT
3300
GTCTGGTGAA ACTAGGGACA AAGTCTACTG ATGAGTTGTC ACTAGGGATC CATTTGATCA
3360
TTTAATCCCA ACAAAAATGA AACAAAATTT TGAGAATTTA TATGCTGAAG TTTTCAACC
3420
CTCTTTTTTA AATAACTTTA TATTATGTAG ATTTGTATTT AGGGTAATTT GTCCAAC TAG
3480
AAGTCCTAAA AATCAATAAA CACACGGATG ACTTTGTCTA ACATTGTATC AGTCATCAAA
3540
TGTAATAATTG TACAAATAAT GAAATTAAAG ATTTAGTCTC TTTTATTTT TTTGTTTAGG
3600
GTGTATATAT ATATATATAT GTATATTGTG TGCATTGATA TATCAATGAG AGGGAGAGAA
3660
CTCAGAGAAG TGTCGGAAAT TAAATGGTA CGAGCCAATT GGAATCTCTG GCATTCTGAG
3720
CTTCATTGTG TTGTTATTAG AAAAAAAAAA AAAAAATCCT TTAAAGATAC CTTCATGATG
3780
ACATTGAATC ATGTAATATA CACGATACAT GGTCTAATTC CTCCTCAAAC CCTAATTACC
3840
AATTCGAAA CCATAATATT TACTAGTATG TTTATATATC CTTACTTTAA GACATTGTTT
3900
GTTTATAATA CCTTGTGAAT TAAGAAAAAA AAAAAAAAC TTGTGGATCT ATTCAAGCCA
3960
TGTGTTAGAA TAAATTTATA AATTTTCTCC TCGTACTGGT CAGATATTGG TCCAACTCC
4020
4080

AAAGCCTTCC CTTTTCAGGA AAAAAACAT TTCGAAATTA ACTCTAATTA ATCAAGAATT
4140
5 TCCTACAATG TATACATCTA ATGTTTTTTC CGCGATCTTA CTTATTAGTG TGAGGGGTAC
4200
AATTGAAAGG TACAAGAAAG CTTGTTCCGA TGCCGTCAAC CCTCCTTCCG TCACCGAAGC
4260
10 TAATACTCAG GTACCAATTT ATATTGTTTG ATTCTCTTTG TTTTATCTTC TTCTTTTCAT
4320
TATATATATG ATCAACAAAA AATATAACCT AAAAAAGAG AGAGTTCAAG GAAATGCATT
4380
15 GAAACGGTTT CGTTATGGTG TTTGAATACA TGGATTTTTG AAGTACTATC AGCAAGAAGC
4440
20 CTCTAAGCTT CGGAGGCAGA TTCGAGATAT TCAGAATTCA AATAGGTAAT TCATTAACTT
4500
TTCATGAAT CTTTCGATTTG GTATTAGGTC ACTTAATTTG GTGTCGGTCC AAAAGTCCGC
4560
25 TTGTAGTTTT CTTTAGAAGT TGTTTTGTTT AATGTTCAATG TTTACAAATT GAAGGCATAT
4620
TGTTGGGGAA TCACTTGGTT CCTTGAACCT CAAGGAACTC AAAACCTAG AAGGACGTCT
4680
30 TGAAAAAGGA ATCAGCCGTG TCCGCTCCAA AAAGGTAAAA TCTACGTTGC TCTCTCTCTG
4740
35 TGTCTCTGTC TCTCTCTCTA TATATAGTCC CTTAGTTTAT ATAGTTCATC ACCCTTTTGT
4800
GAGAATTTTG CAGAATGAGC TGTTAGTGGC AGAGATAGAG TATATGCAGA AGAGGGTAAG
4860
40 AACGTTTCTC CCATTCCAAG TAATTAGATC TTTCTTCGTC TTTGTGAGGG TTTGAGTTTT
4920
CCCATAAATC ATGTGTAGGA AATGGAGTTG CAACACAATA ACATGTACCT GCGAGCAAAG
4980
45 GTTAGCCACG TTCTGTTCCA AATCTTAATC TCAATATCTA CTCTTTTCTT CATTGTATAA
5040
50 CTAAGATAAC GTGAATAACA AGAAAACTTT TGTTTTTGGG TTTAATAGAT AGCCGAAGGC
5100
GCCAGATTGA ATCCGGACCA GCAGGAATCG AGTGTGATAC AAGGGACGAC AGTTTACGAA
5160
55 TCCGGTGTAT CTTCTCATGA CCAGTCGCAG CATTATAATC GGAACATAT TCCGGTGAAC
5220
CTTCTTGAAC CGAATCAGCA ATTCTCCGGC CAAGACCAAC CTCCTCTTCA ACTTGTGTAA
5280
60 CTCAAAACAT GATAACTTGT TTCTTCCCCT CATAACGATT AAGAGAGAGA CGAGAGAGTT
5340
65 CATTTTATAT TTATAACGCG ACTGTGTATT CATAGTTTAG GTTCTAATAA TGATAATAAC
5400
AAAACGTGTG TTTCTTTGCT TAATTACATC AACATTTAAA TCCAAAGTTC TAAAACACGT

5460
CGAGATCCAA AGTTTGT CAT ACAAGATTAG ACGCATACAC GATCAGTTAA TAGATTTTAA
5520
5 GTGCCTTTTA ATATTACAT ATAGTTGCAG CTTGATTAG ATCATGTCCA CCAAACACTC
5580
ACAATTAGAG ACAAGCAAAA CTATAAACAT TGATCATAAA ATGATTACAA CATGTCCATA
10 AATTAATTAT GGATTACAAA AATAAAAACT TACAAAAGAT CT

SEQ ID NO:6 SHP2 genomic

60
15 GAATTCGTAA CAGAATTTAG TGAATAATAT TGTAATTACC AGGCAAGGAC TCTCCAAACG
120
GATAGCTCGA ATATCGTTAT TAAAGAGTAA ATGATCCAAT ATGTAAGCCA TTGTTGATCA
180
20 TCTAACATTG TTGGACTCTC TATTGCTCGA AATGATGCAT ACCTAATCAT TTATTCAGTT
240
AACTATCAAG TTGCATTTGT AAAAACCAAA CATTTAAATT CAGATTTGAT ATCACTTACA
25 300
GAGGATAGAG AAGCATGACT CCAGGCCTGC ATGCAACAAG AAAAAGGAAG AAAATAATGT
360
30 TAAAAATTTG ACAAATATAG TGTTTATTTT TATTATATGA GACAGAATTT GAATAAAATC
420
CTACCCAACT AGAGCATCAA AACGTTTTGC AATCGCAATA ATGAAACCCA TTTTCTTTTT
480
35 GAGTTTTTAC TCTTCTTTCA ACAGAACTT TCTCAAACGT CTTTAGCACT GTGACGTTAG
540
ATATATACAC AAAAGCTTGA AATTTCTTCA AGCAAAAGAA TCTTTGTGGG AGTTAAGGCA
40 600
ACAAGCCAGG TAAAGAATCT CCAACGCATT GTTACGTTTT CATGAACCTA TTTATTATAT
660
45 GTTCTAAGAA AGAAAAAAT ATCTCAAAGT AAACGTTGGA AATTTTCTGA TGAAGGGAAA
720
TCCAAAGTCT TGGGTTTAGT ATCCCTATGA ATGGTATTTG GAATATGTTT TCGTCAAAAC
780
50 AAAAGATTCT TTTCTTTTTC ACAAGAGTTA GTGATCAATA ACTTATGCAC TAATTAATGA
840
GATTGGACGT ATACACAATT TGATTATGAT ACTTGAGTAA AAATCACCTG TCCTTTAATT
900
55 TGGAATCTC TCTTTCTTAC CCATTTATAT ACTACTTCTT TTCATTAAAA TTAAATTTCA
960
60 ATTATCAATC ATCGTTCAAT TTGATAAAGA TTTAACATTT TTTGTCACAG GGCTAGTAAA
1020
AGCAATCTTT ACATAATTCA TCTTTCTTAC ATATATATAT TACCTTTTTTTC TTCATTAGTA
1080
65 TTCTATTGTA TTATGATTAT TTTGTCATAA AGCTAGTAAA TTAAACACTC GATATGAGAA

1140
TTATATTACT TCACGCTAAT TAACTCTTAA CACAACAAGA ACTAGTGCAT ATTCAACTTT
5
1200
CAAAGCATAT ACTATATATT GAGAATATAG ACCACGAAAG TCAATCAAAA GACCTACCAG
1260
CTCTCATCAA GTTCTTTCTT GAAATGATTT TGCAGAATTT CCAACTTAAT TAATTCGACA
10
1320
TGAATGTGAA AATGTGTGTT GCTCGTTAAG AAAATTGAAT AGAAGTACAA TGAAAATGAT
1380
15
GAGGAATGGG CAAAACACAA AAGAGTTTCC TTTCGTAAC ACAAATTAAT AATGCAAATC
1440
TGAGAAAGGG TTCATGGATA ATGACTACAC ACATGATTAG TCATTCCCCG TGGGCTCTCT
20
1500
GCTTTCATTT ACTTTATTAG TTTCATCTTC TCTAATTATA TTGTCGCATA TATGATGCAG
1560
25
TTCTTTTGTC TAAATTACGT AATATGATGT AATTAATTAT CAAAATAAAT ATTCAAATTG
1620
CCGTTGGACT AACCTAATGT CCAAGATTAA GACTTGAACA TAAGAATTTT GGAAAACTA
1680
30
AACCAGTTAT AATATATACT CTAAATTGC CATTCTGAA CACAACCAA TAATAATATA
1740
TACTATTTAC AGTTTTTTTT AATTGGCAAG AACACTGAAA TCTTATTCAT TGTCTCGCTT
35
1800
GGTAGTTGAC AAGTTATAAC ACTCATATTC ATATAACCCC ATTCTAACGT TGACGACGAA
1860
40
CACTCATATA AACCACCCAA ATTCTTAGCA TATTAGCTAA ATATTGGTTT AATTGGAAAT
1920
ATTTTTTTTA TATATAAAAT GCCAGGTAAA TATTAACGAC ATGCAATGTA TATAGGAGTA
45
1980
GGGCAATAAA AAGAAAAGGA GAATAAAAAG GGATTACCAA AAAAGGAAAG TTTCCAAAAG
2040
GTGATTCTGA TGAGAAACAG AGCCCATACC TCTCTTTTTT CCTCTAAACA TGAAAGAAAA
50
2100
ATTGGATGGT CCTCCTTCAA TGCTCTCTCC CCACCCAATC CAAACCCAAC TGTCTTCTTT
2160
55
CTTCTTTTTT TCTTCTTCT AATTTGATAT TTTCTACCAC TTAATTCCAA TCAATTTCAA
2220
ATTTCAATCT AAATGTATGC ATATAGAATT TAATTAAAAG AATTAGGTGT GTGATATTTG
2280
60
AGAAAATGTT AGAAGTAATG GTCCATGTTT TTTCTTTCTT TTTCTTCTA TAACACTTCA
2340
GTTTGAAAAA AAACCTACCAA ACCTTCTGTT TTCTGCAAT GGGTTTTTAA ATACTTCCAA
65
2400
AGAAATATTC CTCTAAAAGA AATTATAAAC CAAAACAGAA ACCAAAAACA AAAAATAAAG
2460

TTGAAGCAGC AGTTAAGTGG TACTGAGATA ATAAGAATAG TATCTTTAGG CCAATGAACA
2520
5 AATTAAGTCT CTCATAATTC ATCTTCCCAT CCTCACTTCT CTTTCTTTCT GATATAATTA
2580
ATCTTGCTAA GCCAGGTATG GTTATTGATG ATTTACACTT TTTTAAAAA GTTTCCTCCT
2640
10 TTTCTCCAAT CAAATTCTTC AGTTAATCCT TATAAACCAT TTCTTTAATC CAAGGTGTTT
2700
GAGTGCAAAA GGATTTGATC TATTTCTCTT GTGTTTATAC TTCAGCTAGG GCTTATAGAA
2760
15 ATGGAGGGTG GTCCGAGTAA TGAAGTAGCA GAGAGCAGCA AGAAGATAGG GAGAGGGAAG
2820
20 ATAGAGATAA AGAGGATAGA GAACACTACG AATCGTCAAG TCACTTTCTG CAAACGACGC
2880
AATGGTTTAC TCAAGAAAGC TTATGAGCTC TCTGTCTTGT GTGACGCTGA GGTGCTCTT
2940
25 GTCATCTTCT CCACTCGAGG CCGTCTCTAC GAGTACGCCA ACAACAGGTA CACATCTTTT
3000
AGCTAGATCT TGATTTTGTG GAATTTTTTT TCTAGAATAA AGTTTCGACT CTTCTGGTGG
3060
30 GTTTTCAAT CTTTATGGTC TCTTATAGT TTTTTCCTT AGTTTCTCTG AAGCTCAAAT
3120
35 CTCTTTAAAA ATCCCCAAAA TTAGGGTTTG TTTAAACTA GGAACCCCTA CTTTAACTTC
3180
TTTCTCTAG TAAAAAAGCA GTGAGGGTCT TCTCTGATCA TTAATTAGCA TCCCCATAC
3240
40 CTTGTTCCAG TCACTTTTTC TCCACAAATC CTTATAACAG TATCTATATA TGTATCTATT
3300
TATGTCAGTT TGTACAAGAC ACTTCGATCA ATTTGATGAC CCATCAAGTT TTATTCTGCT
3360
45 AGATTGATCA TTAGGTTTCC ATCATAGTAA TGAAAAAGTA GGGTCTTGA TAAATTATA
3420
50 ATAATATATA TTATTTGGCT ATATAAAAAA GCTATGTAGA TTCCTTAAAA ATTGATTCAC
3480
TAGGGAGAGA CTAGTAGGTG TTTGTCTTCT GACACTTCTC TAATCTTTTG GTGAATCCTT
3540
55 TTGTTAAATC AAGAAAATGA ATCAGGGACA AAGCTTATTG TTGAGTCACT TAATTAATCA
3600
TCCGATCCAT CAATCAAGAA AAATAACGAA ACAGAAAATT TTGATTTTTC ATTGTTATTT
3660
60 TCTCCACTTC AAGTTGGGGA CTTGTCATTT CCGTTTTTCT ATACGTTTCC AGCTATTAAAC
3720
65 AGCTCATGTT CATTTCACCA TTTTGATTAT TTGTCTGCTT TTTAAAGATA AATGTTTTCA
2780
AAAATATTGT TTTTATTTGC TTGGCTAGTT AATACTATAA TTGAGGTGTA TGTATGACTA

3840
TAATCTATAA GTCAAGTCTC ATATCATGGA TCTAAGTTAA AACTAGTAAA TTTGTAGTTT

5 3900
CAATGTGAAC TTTCACAACG ACTAAAGAAC TGATCTGAAG TTTATAATGG ACATGACTAA

3960
TTTGATTAAC AAAAGAGGAA TGCATTATGT ATGTAGAAAC ATGTGATATA TATATGTTTC

10 4020
TATTATCAAA AGTGTAGTTA ACTTTCTTAT TTCAAACACC CTCATGCTTT AGTAGTATCT

4080
TACTTTTGAC ATTTCTCAAC TTCAGCTTTC CATTATACAA CAGCACAATG TAAATTACTT

15 4140
GTATATGAAT ATGAAAGCAT AACGTTATGC AAAGATTCTT AGCTTTTCTT TTTCTGTTTT

4200
GCAAAAGATT TACAAATATC ATGTTCTTGG TAAAAACATA CTGCGCTCAG CCACATATGC

20 4260
ATGTAAATGT AATGTTCAAA TATTAATTCA GGAAAAACAA AGAAGAAGCA AAATTAGCTT

4320
CTAGAGTAGG GAATCTATTG ACTTGACCTG AAAATCACTT CTTTTTCTTA AAGCCTAGTA

4380
GTGAATTTTT TAATCTAATT AGGCCAAAAT ATATACTAGC CTAAAATATA ATTTGGATTT

30 4440
TGTGTCGTAC ATAAATTGGG ACCAATTCCA ATTAATAAG AGCATATGCA ATTCAAATTC

4500
TTTTTATTTT CTCTCCGAT TTGCTACTTC TTTCTTTTGT ATGTTTTCAA ATTAGGATTA

35 4560
CACTTTTTTG GGAAGTACA CATTAGGGTC TTCTCGAACT TTGATTATAC ATATATATAT

4620
ATATATATAT ATATACTTT GTGAGATGTC ACTGTTAATA GATAATAGGC AATAACAATA

4680
ATATCCAAA AAGAAGGCGC AAACAAATCA TATACTATAT GGTACTGGTC CATTCACTAT

45 4740
TTTGTCGGTT GAATTTAAGG TTGGCGTAC AAACTTTGTT TCAAACCTTT ATTATTCCGT

4800
CTTTCTGTGT GTTTTGTATA TCCAGAAGAT AAAAATATCA ATTTCTTTAA CGACTTCATA

50 4860
TATATATATA TATATATATA TATATATATT TTTCTCTTCT GGTTTTAGTG TTTGAATCCA

4920
ACAGTTATAG TTTCGTGTGT CTTTGTTTTA CTTGTGGTGG TTTAAGTTTG AGATTTTCAC

4980
CGATTGCATC TATTTACATA TATAGCTACC ACAAAAAGA TTGCATTTTA AAATCTTTTC

60 5040
CTTTGTGTGA ATGTTGATGA AGTGTGAGAG GAACAATAGA AAGGTACAAG AAAGCTTGCT

5100
CCGACGCCGT TAACCCTCCG ACCATCACCG AAGCTAATAC TCAGGTTAGC TTTTAATTAA

65 5160
TACACCTAGC TAGCTAGTTC GTTAATTACT TAATTTCTTC TTCTTTTAGT TATCTGACCT

5220
TTTTTTCACC TCTTGTAACA ATGATGGGAT CGAAATTGAT GAAGTACTAT CAGCAAGAGG
5
5280
CGTCTAAACT CCGGAGACAG ATTCGGGACA TTCAGAATTT GAACAGACAC ATTCTTGGTG
5340
AATCTCTTGG TTCCTTGAAC TTTAAGGAAC TCAAGAACCT TGAAAGTAGG CTTGAGAAAG
10
5400
GAATCAGTCG TGTCCGATCC AAGAAGGTAC ATCACTAACT CTCCATCAAT CTCCTTATCA
5460
15
TTGAATATAT ATCCATCTGA TTCTTGCCCG TTATATTTGG TTTTCTCTC CAGCAGAGA
5520
TGTTAGTTGC AGAGATTGAA TACATGCAAA AAAGGGTAAA AGTAAACCT ATCTTCCTTC
20
5580
ACAATGAACT ACCCCTACTT TATTAGCAAC TTCTCTTTCT GATGATCATC TTTTATTAT
5640
25
TCTGTTGTCG CTGCAATTGT AGGAAATCGA GCTGCAAAAC GATAACATGT ATCTCCGCTC
5700
CAAGGTTTTA TACATAACTC TTTTGGCAT TTTTGATCAT CATTTTTTTC CGGTAGACAA
5760
30
TCTCTTGATG TGCAAATCT AAATATCTCT GCAGATTACT GAAAGAACAG GTCTACAGCA
5820
ACAAGAATCG AGTGTGATAC ATCAAGGGAC AGTTTACGAG TCGGGTGTTA CTTCTTCTCA
35
5880
CCAGTCGGGG CAGTATAACC GGAATTATAT TCGGTTAAC CTTCTTGAAC CGAATCAGAA
5940
40
TTCCTCCAAC CAAGACCAAC CACCTCTGCA ACTTGTTTGA TTCAGTCTAA CATAAGCTTC
6000
TTTCCTCAGC CTGAGATCGA TCTATAGTGT CACCTAAATG CGGCCGCGTC CCTCAACATC
6060
45
TAGTCGCAAG CTGAGGGGAA CCACTAGTGT CATACGAACC TCCAAGAGAC GGTACACAA
6120
ACGGGTACAT TGTGATGTC ATGTATGACA ATCGCCCAAG TAAGTATCCA GCTGTGTTCA
50
GAACGTACGT CCGAATTC